

BIRCH, STEWART, KOLASCH & BIRCH, LLP

INTELLECTUAL PROPERTY LAW
8110 GATEHOUSE ROAD
SUITE 500 EAST
FALLS CHURCH, VA 22042-1210
U S A
(703) 205-8000

FAX: (703) 205-8050
(703) 698-8590 (G IV)

e-mail: mailroom@bskb.com
web: <http://www.bskb.com>

CALIFORNIA OFFICE:
COSTA MESA, CALIFORNIA

THOMAS S. AUCHTERLONIE
MICHAEL R. CAMMARATA
JAMES T. ELLER, JR.
SCOTT L. LOWE
MARK J. NUEL, Ph.D.
DARIN E. BARTHOLOMEW*
D. RICHARD ANDERSON
PAUL C. LEWIS
W. KARL RENNER
MARK W. MILSTEAD*
JOHN CAMPA*
REG. PATENT AGENTS
FREDERICK R. HANDREN
ANDREW J. TELESZ, JR.
MARYANNE ARMSTRONG, Ph.D.
MAKI HATSUMI
MIKE S. RYU
CRAIG A. McROBBIE
GARTH M. DAHLEN, Ph.D.
LAURA C. LUTZ
ROBERT E. GOOZNER, Ph.D.
HYUNG N. SOHN
MATTHEW J. LATTIG
ALAN PEDERSEN-GILES
JUSTIN D. KARJALA

TERRELL C. BIRCH
RAYMOND C. STEWART
JOSEPH A. KOLASCH
JAMES M. SLATTERY
BERNARD L. SWEENEY*
MICHAEL K. MUTTER
CHARLES GORENSTEIN
GERALD M. MURPHY, JR.
LEONARD R. SVENSSON
PERRY L. CLARK
ANDREW D. MEIKLE
MARC S. WEINER
JOE MCKINNEY MUNCY
ROBERT J. KENNEY
DONALD J. DALEY
JOHN W. BAILEY
JOHN A. CASTELLANO, III
GARY D. YACURA
OF COUNSEL
HERBERT M. BIRCH (1905-1996)
ELLIOT A. GOLDBERG*
WILLIAM L. GATES*
EDWARD H. VALANCE
RUPERT J. BRADY (RET.)*

*ADMITTED TO A BAR OTHER THAN VA

Date: January 27, 2000
(First day after Government Snow Closure)

Docket No.: 2815-127P

BOX PATENT APPLICATION

Assistant Commissioner for Patents
Washington, DC 20231

Sir:

As authorized by the inventor(s), transmitted herewith for filing is a patent application applied for on behalf of the inventor(s) according to the provisions of 37 C.F.R. § 1.41(c), which claims priority under 35 U.S.C. § 119(e) of Provisional Application No. 60/118,112 filed on February 1, 1999

Inventor(s): Thomas J. JENTSCH

For: NOVEL POTASSIUM CHANNELS AND GENES ENCODING THESE POTASSIUM CHANNELS

Enclosed are:

- ☒ A specification consisting of THIRTY-SEVEN (37) pages plus a Sequence Listing (11 pages)
- ☒ Four (4) sheet(s) of formal drawings
- ☐ Certified copy of Priority Document(s)
- ☒ Executed Declaration in accordance with 37 C.F.R. § 1.64 will follow
- ☐ A statement to establish small entity status under 37 C.F.R. § 1.9 and 37 C.F.R. § 1.27
- ☒ Preliminary Amendment

- ✓ ☒ Information Sheet
- ☐ Information Disclosure Statement, PTO-1449 and reference(s)
- ☒ Amend the specification by inserting before the first line the sentence:

--This application claims priority on provisional Application No. 60/118,112 filed on February 1, 1999, the entire contents of which are hereby incorporated by reference.--

- ☐ Other:

The filing fee has been calculated as shown below:

			LARGE ENTITY	SMALL ENTITY	
BASIC FEE			\$690.00	\$345.00	
	NUMBER FILED	NUMBER EXTRA	RATE FEE	RATE FEE	
TOTAL CLAIMS	56- 20 =	36	X 18 = \$648.00	x 9 = \$0.00	
INDEPENDENT CLAIMS	6- 3 =	3	x 78 = \$234.00	x 39 = \$0.00	
<input type="checkbox"/> MULTIPLE DEPENDENT CLAIMS PRESENTED			+ \$260.00	+ \$130.00	
			TOTAL	\$1,572.00	\$0.00

- ☒ The application transmitted herewith is filed in accordance with 37 C.F.R. § 1.41(c). The undersigned has been authorized by the inventor(s) to file the present application. The original duly executed declaration together with the surcharge will be forwarded in due course.
- ✓ ☒ A check in the amount of \$1,572.00 to cover the filing fee is enclosed.
- ☐ Please charge Deposit Account No. 02-2448 in the amount of \$0.00. A triplicate copy of this transmittal form is enclosed.

☒ Please send correspondence to:

BIRCH, STEWART, KOLASCH & BIRCH, LLP **or** Customer No. 2292
P.O. Box 747
Falls Church, VA 22040-0747
Telephone: (703) 205-8000

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. §§ 1.16 or 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By 
Gerald M. Murphy, Jr., #28,977

P.O. Box 747
Falls Church, VA 22040-0747
(703) 205-8000

GMM/afy
2815-127P

Attachments

(Rev. 01/08/2000)

PATENT
2815-127P

IN THE U.S. PATENT AND TRADEMARK OFFICE

Applicants: Thomas J. JENTSCH
Serial No.: NEW Group:
Filed: January 27, 2000 Examiner:
For: NOVEL POTASSIUM CHANNELS AND GENES ENCODING THESE
POTASSIUM CHANNELS

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

January 27, 2000
(First day after Gov't
Snow Closure)

Sir:

The following preliminary amendments and remarks are respectfully submitted in connection with the above-identified application.

IN THE CLAIMS

Please cancel claims 46 and 48 without prejudice or disclaimer of the subject matter contained therein.

Please amend the following new claims:

Claim 1 (Amended)

Line 2, change "hybridising" to --hybridizing--

Line 3, change "presented as" to --of--

Claim 2 (Amended) The isolated polynucleotide according to claim 1, [being] wherein said isolated polynucleotide is at least 50% homologous, preferably more than 70% homologous, more [preferred] preferably more than 80% homologous, even more [preferred] preferably more than 90% homologous, and most [preferred] preferably more than 95%[,] homologous to the polynucleotide sequence [presented as] of SEQ ID NO: 1.

Claim 3 (Amended) The isolated polynucleotide according to [either of claims 1-2 being] claim 1, wherein said isolated polynucleotide is a cloned polynucleotide.

Claim 4 (Amended) The isolated polynucleotide according to claim 3, [in which] wherein the polynucleotide is cloned from, or produced [on the basis of] from a cDNA library.

Claim 5 (Amended)

Line 1, change "any of claims 1-4" to --claim 1--

Line 2, change "presented as" to --of--

Line 3, change "hereof" to --thereof--

Claim 6 (Amended) The isolated polynucleotide according to [any of claims 1-4] claim 1, comprising the polynucleotide sequence [presented as] of SEQ ID NO: 1, or a sub-sequence [hereof, including the] thereof, wherein said sequence includes a mutation G935A.

Claim 7 (Amended) The isolated polynucleotide according to [any of claim 1-6, encoding] claim 1, wherein said isolated polynucleotide encodes a potassium channel, or a potassium channel subunit.

Claim 8 (Amended)

Line 1, change "the" (second occurrence) to --a--

Line 2, change "represented by" to --of--

Line 3, change "hereof" to --thereof--

Claim 9 (Amended)

Line 2, change "which" to --wherein said--

Claim 10 (Amended) The isolated polynucleotide according to claim 9, [which] wherein said variant has an amino acid sequence that has been changed at one or more positions located in [the] a conserved [regions, as] region, wherein said region is defined by Table 1.

Claim 11 (Amended) The isolated polynucleotide according to claim 9, encoding [the] a variant KCNQ4/G285S [(i.e. KCNQ4/G333S according to the KCNQ1 numbering)] or KCNQ4/G333S when said polynucleotide is numbered according to KCNQ1.

Claim 12 (Amended)

Line 1, change "set forth in" to --of--

Claim 13 (Amended)

Line 2, change "claims 1-11" to --claim 1--

Claim 14 (Amended) The polypeptide according to claim 13, [being] wherein said polypeptide is a KCNQ4 potassium channel subunit comprising the amino acid sequence [presented as] of SEQ ID No. 2.

Claim 15 (Amended) The polypeptide according to claim 13, [being a] wherein said polypeptide is a KCNQ4 variant, [which] and wherein said variant has an amino acid sequence that has been changed by deletion of an amino acid residue, by insertion of an additional amino acid residue, or by substitution of an amino acid residue at one or more positions.

Claim 16 (Amended) The polypeptide according to claim 15, [which] wherein said variant has an amino acid sequence that has been changed at one or more positions located in [the] a conserved [regions, as] region, wherein said region is defined by Table 1.

Claim 17 (Amended) The polypeptide according to claim 15, [being] wherein said polypeptide is the variant KCNQ4/G285S [(i.e. KCNQ4/G333S according to the KCNQ1 numbering)] or KCNQ4/G333S when said polypeptide is numbered according to KCNQ1.

Claim 18 (Amended)

Line 2, change "any of claims 1-11" to --claim 1--

Claim 19 (Amended)

Line 2, change "presented as" to --of--

Line 3, change "hereof" to --thereof--

Claim 20 (Amended)

Line 2, change "which" to --wherein said--

Claim 21 (Amended) The cell according to claim 20, [which] wherein said variant has an amino acid sequence that has been changed at one or more positions located in [the] a conserved [regions, as] region, wherein said region is defined by Table 1.

Claim 22 (Amended) The cell according to claim 20, genetically manipulated by the incorporation of the variant KCNQ4/G285S [(i.e. KCNQ4/G333S according to the KCNQ1 numbering)] or KCNQ4/G333S when numbered according to KCNQ1.

Claim 23 (Amended) The cell according to [any of claims 18-22, genetically manipulated to co-express] claim 18, wherein said cell co-expresses one or more KCNQ channel subunits.

Claim 24 (Amended) The cell according to 23, [genetically manipulated to co-express] wherein said cell co-expresses KCNQ4 and KCNQ1 channel subunits[;], KCNQ4 and KCNQ2 channel subunits[;], KCNQ4 and KCNQ3 channel subunits[;], KCNQ4 and KCNQ1 and KCNQ2 channel subunits[;], KCNQ4 and KCNQ1 and KCNQ3 channel

subunits[;] KCNQ4 and KCNQ2 and KCNQ3 channel subunits[;] or KCNQ4 and KCNQ1 and KCNQ2 and KCNQ3 channel subunits.

Claim 25 (Amended) The cell according to claim 23, [genetically manipulated to co-express] wherein said cell co-expresses KCNQ3 and KCNQ4 channel subunits.

Claim 26 (Amended) The cell according to [any of claims 18-25, being] claim 18, wherein said cell is an eukaryotic cell, in particular a mammalian cell, an oocyte, or a yeast cell.

Claim 27 (Amended) The cell according to [any] claim 26, [being] wherein said cell is a human embryonic kidney (HEK) cell, a HEK 293 cell, a BHK21 cell, a Chinese hamster ovary (CHO) cell, a *Xenopus laevis* oocyte (XLO) cell, a COS cell, or any other cell line [able to express] that expresses KCNQ potassium channels.

Claim 28 (Amended)

Line 1, change "any of claims 18-27" to --claim 18--

Claim 29 (Amended) A method for obtaining a substantially homogeneous source of a human potassium channel[,] comprising a KCNQ4 subunit, [which method comprises] comprising the steps of:
culturing a cellular host having incorporated expressibly therein a polynucleotide according to [any of claims 1-11, and then] claim 1; and
recovering the cultured cells.

Claim 30 (Amended)

Line 1, before "comprising" insert --further-- and delete "subsequent"

Claim 31 (Amended) A method of [screening] detecting a chemical compound [for capability of binding] which binds to a potassium channel comprising at least one KCNQ4 channel subunit[, which method comprises] comprising the steps of:

(i) subjecting a KCNQ4 channel subunit [containing] of a cell [according to claims 18-27,] or a membrane preparation [according to claim 28,] to the action of a KCNQ4 binding agent to form a complex [with the KCNQ4 channel subunit containing cell];

(ii) subjecting the complex of step (i) to the action of the chemical compound to be tested; and

(iii) detecting the displacement of the KCNQ4 binding agent from the complex with the KCNQ4 channel subunit [containing] of the cell or membrane preparation.

Claim 32 (Amended) The method of claim 31, wherein the cell containing the KCNQ4 channel subunit [containing cell] is a cell, [according to any of claims 18-27, or a membrane preparation according to claim 28] which is genetically manipulated by the incorporation of a heterologous polynucleotide having a nucleic acid sequence which is capable of hybridizing under high

stringency conditions with the polynucleotide of SEQ ID NO. 1, its
complimentary strand, or a sub-sequence thereof.

Claim 33 (Amended)

Line 1, change "either of claims 31-32, in which" to --claim
31, wherein--

Line 3, change "(Linopirdine);" to --(Linopirdine),--

Claim 34 (Amended) The method of claim 33, [which compounds
have been marked] wherein said binding agents are radioactively
labelled with ^3H .

Claim 35 (Amended)

Line 1, change "either of claims 33-34" to --claim 33--

Line 2, delete "containing"

Line 3, change "the" to --an--

Claim 36 (Amended) A method of screening a chemical compound
for activity on a potassium channel comprising at least one KCNQ4
channel subunit, [which method comprises] comprising the steps of:

(i) subjecting a KCNQ4 channel subunit [containing] of a cell
to the action of the chemical compound; and

(ii) monitoring the membrane potential, the current, the
potassium flux, or the secondary calcium influx of the KCNQ4
channel subunit [containing] of said cell.

Claim 37 (Amended)

Line 1, change "the KCNQ4 channel subunit containing" to --
said--

Line 2, change "any of claims 18-27" to --claim 18--

Claim 38 (Amended)

Line 1, change "either of claims 36-37" to --claim 36--

Line 2, delete "of the KCNQ4 channel subunit containing cell"

Claim 39 (Amended)

Line 1, change "either of claims 36-37" to --claim 36--

Line 2, delete "of the KCNQ4 channel subunit containing cell"
and change "using" to --by--

Claim 40 (Amended)

Line 1, change "claims 31-35, and/or by the" to --claim 31--

Line 2, delete "method of claims 36-39"

Claim 41 (Amended) [Use of the chemical compound according to
claim 40 for] A method of diagnosis, treatment, prevention or
alleviation of diseases related to tinnitus, loss of hearing, [in
particular] progressive hearing loss, neonatal deafness, and
presbycusis, [(deafness of the elderly); and] diseases of adverse
conditions of the CNS, [including] affective disorders,
Alzheimer's disease, anxiety, ataxia, CNS damage caused by trauma,
stroke or neurodegenerative illness, cognitive deficits,

compulsive [behaviour] behavior, dementia, depression, Huntington's disease, mania, memory impairment, memory disorders, memory dysfunction, motion disorders, motor disorders, neurodegenerative diseases, Parkinson's disease and Parkinson-like motor disorders, phobias, Pick's disease, psychosis, schizophrenia, spinal cord damage, stroke, and tremor comprising the step of:

using a chemical compound of claim 40.

Claim 42 (Amended)

Line 1, change "use" to --method--

Line 2, change "(Linopirdine);" to --(Linopirdine),--

Claim 43 (Amended) [Use of a polynucleotide sequence according to any of claims 1-12, for the] A method for screening [of] genetic materials [for] of individuals [having this mutations] comprising:

contacting a polynucleotide of claim 1 to said genetic material; and

detecting hydridization of said polynucleotide to said genetic material.

Claim 44 (Amended)

Line 2, change "gene, a" to --gene or a-- and change "gene, or genetically manipulated in order to over-" to --gene.--

Line 3, delete in its entirety.

Claim 45 (Amended)

The transgenic animal according to claim 44, [being a knock-out animal in which the gene is totally deleted] wherein said knock-out is in a homozygous state.

Claim 47 (Amended) The transgenic animal according to [any of claims 44-46, being] claim 44, wherein said transgenic animal is a transgenic rodent, in particular a hamster, a guinea pig, a rabbit, or a rat, a transgenic pig, a transgenic cattle, a transgenic sheep, or a transgenic goat.

Claim 49 (Amended) [The use according to claim 48, for the screening of] A method to screen for drugs affecting diseases or conditions associated with hearing loss or tinnitus, comprising injecting a transgenic animal of claim 44 with a therapeutic compound.

Claim 50 (Amended) A method for [the identification, localization, isolation or amplification] identifying, localizing, isolating or amplifying a polynucleotide [according to any one of claim 1-11], comprising using a polynucleotide according to claim 12 as a primer or a probe.

Claim 51 (Amended) An antibody capable of binding one or more polypeptides [as claimed in any one or claims 13-17] according to claim 13.

Claim 52 (Amended) The antibody of claim 51, [being a] wherein said antibody is a monoclonal antibody.

Please add the following new claims:

--53. The method of claim 31, wherein a membrane preparation is used.

54. A chemical compound identified by the method of claim 36.

55. A method of diagnosis, treatment, prevention or alleviation of diseases related to tinnitus, loss of hearing, progressive hearing loss, neonatal deafness, and presbycusis, diseases of adverse conditions of the CNS, affective disorders, Alzheimer's disease, anxiety, ataxia, CNS damage caused by trauma, stroke or neurodegenerative illness, cognitive deficits, compulsive behavior, dementia, depression, Huntington's disease, mania, memory impairment, memory disorders, memory dysfunction, motion disorders, motor disorders, neurodegenerative diseases, Parkinson's disease and Parkinson-like motor disorders, phobias, Pick's disease, psychosis, schizophrenia, spinal cord damage, stroke, and tremor comprising the step of:

using a chemical compound of claim 54.

56. The method according to claim 55, wherein the chemical compound is 1,3-dihydro-1-phenyl-3,3-bis(4-pyridylmethyl)-2H-indol-2-one (Linopirdine), or 10-10-bis(4-pyridinyl-methyl)-9(10H)-anthracenone.

57. A transgenic animal that overexpresses a KCNQ4 gene or a mutated KCNQ4 gene.

58. The transgenic animal according to claim 57, wherein said transgenic animal is a transgenic rodent, in particular a hamster, a guinea pig, a rabbit, or a rat, a transgenic pig, a transgenic cattle, a transgenic sheep, or a transgenic goat.--


REMARKS

Entry of the above amendments is earnestly solicited. An early and favorable first action on the merits is earnestly solicited.

If necessary, the Commissioner is hereby authorized in this, concurrent, and further replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fee required under 37 C.F.R. 1.16 or under 37 C.F.R. 1.17; particularly, extension of time fees.

Respectfully yours,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By 
Gerald M. Murphy, Jr.
Reg. No. 28,977

P.O. Box 747
Falls Church, VA 22048-0747
(703) 205-8000

GMM/KLR/afy

NOVEL POTASSIUM CHANNELS AND GENES ENCODING THESE POTASSIUM CHANNELS

TECHNICAL FIELD

5

This invention relates to novel potassium channels and genes encoding these channels. More specifically the invention provides isolated polynucleotides encoding the KCNQ4 potassium channel subunit, cells transformed with these polynucleotides, transgenic animals comprising genetic mutations, and the use of the transformed cells and the transgenic animals for the *in vitro* and *in vivo* screening of chemical compounds affecting KCNQ4 subunit containing potassium channels.

BACKGROUND ART

15

Potassium channels participate in the regulation of electrical signalling in excitable cells, and regulates the ionic composition of biological fluids. Mutations in the three known genes of the KCNQ branch of the K⁺-channel gene family underlie inherited cardiac arrhythmia's, in some cases associated with deafness, and neonatal epilepsy.

20

Hearing loss is the most frequent sensory defect in humans. Hearing loss can be due to environmental and genetic factors, and the progressive hearing loss of the elderly (presbycusis) most often seems to be due to a combination of both.

Inherited deafness can be classified as non-syndromic (isolated hearing loss) or syndromic (associated with other anomalies). Several hundred syndromes, consisting of hearing loss associated with defects in a variety of other organ systems, have been described. Non-syndromic deafness is classified according to its mode of inheritance as DFN, DFNA, and DFNB (X-linked, autosomal dominant and autosomal recessive, respectively). In general, autosomal recessive deafness has an early onset and is very severe. Autosomal dominant deafness, by contrast, more often develops slowly over several decades and may become apparent only in adulthood. It is hoped that genes identified in families with dominant deafness may also - with different types of mutations - underlie some forms of presbycusis.

A bewildering number of loci for non-syndromic deafness were identified in the last four years. There are at least 19 loci for autosomal dominant deafness (DFNA1 to DFNA19), and 22 loci for DFNB. Sometimes, depending on the particular mutation, the same gene can be involved in dominant or recessive deafness. This large number of loci reflects the complexity of the inner ear. Identification of these genes and characterisation of their products will significantly advance our understanding of the molecular basis of the physiology of this sensory organ.

Several genes involved in syndromic and non-syndromic deafness have already been identified and are reviewed by *Petit* [*Petit C*: Genes responsible for human hereditary deafness: symphony of a thousand; *Nature Genet.* 1996 **14** 385-391] and *Kalatzis & Petit* [*Kalatzis V & Petit C*: The fundamental and medical impacts of recent progress in research on hereditary hearing loss; *Hum. Mol. Genet.* 1998 **7** 1589-1597]. Among others, their gene products include transcription factors, unconventional myosin isoforms, α -tectorin (an extracellular matrix protein), diaphanous, a protein interacting with the cytoskeleton, connexin 26 (a gap junction protein), and two genes encoding potassium channel subunits, *KCNQ1* and *KCNE1*.

Ion channels play important roles in signal transduction and in the regulation of the ionic composition of intra- and extracellular fluids. Mutations in ion channels were since long suspected as possibly underlying some forms of hearing loss. In the cochlea (the auditory sensory organ), the transduction current through the sensory cells is carried by potassium ions and depends on the high concentration of that ion in the endolymph. So far only two genes encoding potassium channel subunits, *KCNQ1* and *KCNE1*, were found to be mutated in syndromic hereditary deafness. The gene products of both genes, the *KCNQ1* (or *KvLQT1*) and the *minK* (or *IsK*) protein, respectively, form heteromeric potassium channels.

KCNQ1 is a typical member of the voltage-gated potassium channel superfamily with 6 transmembrane domains and a pore region situated between the fifth and the sixth transmembrane domain. The *minK* protein has a single transmembrane span and cannot form potassium channels on its own. However, as a β -subunit it enhances and modifies currents mediated by *KCNQ1*. These heteromeric channels participate in the repolarization of the heart action potential. Certain mutations in either *KCNQ1* or *KCNE1* cause a form of the autosomal dominant long QT syndrome (LQTS), a disease characterised by repolarization anomalies of cardiac action potentials resulting in arrhythmias and sudden death. Interestingly, other mutations in either gene lead to the recessive Jervell and Lange-Nielsen (JLN) syndrome that combines LQTS with congenital deafness. In order to cause deafness, *KCNQ1/minK* currents must be reduced below levels that are already sufficiently low to cause cardiac arrhythmia.

SUMMARY OF THE INVENTION

We have now cloned and characterised *KCNQ4*, a novel member of the *KCNQ* family of potassium channel proteins. *KCNQ4* has been mapped to the DFNA2 locus for autosomal dominant hearing loss, and a dominant negative *KCNQ4* mutation that causes deafness in a DFNA2 pedigree was identified.

KCNQ4 is the first potassium channel gene underlying non-syndromic deafness. *KCNQ4* forms heteromeric channels with other *KCNQ* channel subunits, in particular *KCNQ3*.

The present invention has important implications for the characterisation and exploitation of this interesting branch of the potassium channel super family, as well as for the understanding of the cochlear physiology, and for human deafness and progressive hearing loss.

Accordingly, in its first aspect, the invention provides an isolated polynucleotide having a nucleic acid sequence which is capable of hybridising under high stringency conditions with the polynucleotide sequence presented as SEQ ID NO: 1, its complementary strand, or a sub-sequence thereof.

In another aspect the invention provides a recombinantly produced polypeptide encoded by the polynucleotide of the invention.

In a third aspect the invention provides a cell genetically manipulated by the incorporation of a heterologous polynucleotide of the invention.

In a fourth aspect the invention provides a method of screening a chemical compound for inhibiting or activating or otherwise modulating the activity on a potassium channel comprising at least one *KCNQ4* channel subunit, which method comprises the steps of subjecting a *KCNQ4* channel subunit containing cell to the action of the chemical compound; and monitoring the membrane potential, the current, the potassium flux, or the secondary calcium influx of the *KCNQ4* channel subunit containing cell.

In a fifth aspect the invention relates to the use of a polynucleotide sequence of the invention for the screening of genetic materials from humans suffering from loss of hearing (e.g. dominant, recessive, or otherwise), tinnitus, and other neurological diseases for mutations in the *KCNQ4* gene.

In a sixth aspect the invention relates to the chemical compound identified by the method of the invention, in particular to the use of such compounds for diagnosis, treatment or alleviation of a disease related to tinnitus; loss of hearing, in particular progressive hearing loss, neonatal deafness, and presbycusis (deafness of the elderly); and diseases or adverse conditions of the CNS, including affective disorders, Alzheimer's disease, anxiety, ataxia, CNS damage caused by trauma, stroke or neurodegenerative illness, cognitive deficits, compulsive behaviour, dementia, depression, Huntington's disease, mania, memory impairment, memory disorders, memory dysfunction, motion disorders, motor disorders, neurodegenerative diseases, Parkinson's disease and Parkinson-like motor disorders, phobias, Pick's disease, psychosis, schizophrenia, spinal cord damage, stroke, and tremor.

In a seventh aspect the invention provides a transgenic animal comprising a knock-out mutation of the endogenous *KCNQ4* gene, a replacement by or an additional expression of a mutated *KCNQ4* gene, or genetically manipulated in order to over-express the *KCNQ4* gene or to over-express mutated *KCNQ4* gene.

- 5 In an eighth aspect the invention relates to the use of the transgenic animal of the invention for the *in vivo* screening of therapeutic compounds.

Other objects of the invention will be apparent to the person skilled in the art from the following detailed description and examples.

10 DETAILED DISCLOSURE OF THE INVENTION

- The present invention provides novel voltage-gated potassium channels and genes encoding these channels. The invention also provides cells transformed with these genes, transgenic animals comprising genetic mutations, and the use of the
15 transformed cells and the transgenic animals for the *in vitro* and *in vivo* screening of drugs affecting *KCNQ4* containing potassium channels.

Polynucleotides

In its first aspect, the invention provides novel polynucleotides.

- 20 The polynucleotides of the invention are such which have a nucleic acid sequence capable of hybridising under high stringency conditions with the polynucleotide sequence presented as SEQ ID NO: 1, its complementary strand, or a sub-sequence thereof.

- The polynucleotides of the invention include DNA, cDNA and RNA
25 sequences, as well as anti-sense sequences, and include naturally occurring, synthetic, and intentionally manipulated polynucleotides. The polynucleotides of the invention also include sequences that are degenerate as a result of the genetic code.

- As defined herein, the term "polynucleotide" refers to a polymeric form of nucleotides of at least 10 bases in length, preferably at least 15 bases in length. By
30 "isolated polynucleotide" is meant a polynucleotide that is not immediately contiguous with both of the coding sequences with which it is immediately contiguous (one on the 5' end and one on the 3' end) in the naturally occurring genome of the organism from which it is derived. The term therefore includes recombinant DNA which is incorporated into an expression vector, into an autonomously replicating plasmid or virus, or into the
35 genomic DNA of a prokaryote or eukaryote, or which exists as a separate molecule, e.g. a cDNA, independent from other sequences.

The polynucleotides of the invention also include allelic variants and "mutated polynucleotides" comprising a nucleotide sequence that differs from the

sequence presented as SEQ ID NO: 1 at one or more nucleotide positions. The mutated polynucleotide may in particular be a polynucleotide of the invention comprising a nucleotide sequence as in SEQ ID NO: 1, which sequence, however, differs from SEQ ID NO: 1 so as to effect the expression of a variant polypeptide. The mutated polynucleotide may be a polynucleotide of the invention having a nucleotide sequence encoding a potassium channel having an amino acid sequence that has been changed at one or more positions. The mutated polynucleotide may in particular be a polynucleotide of the invention having a nucleotide sequence encoding a potassium channel having an amino acid sequence that has been changed at one or more positions located in the conserved regions, as defined by Table 1, below.

In a more specific embodiment the polynucleotide of the invention has the polynucleotide sequence giving rise to the G285S mutation as indicated in SEQ ID NO: 1, i.e. the DNA sequence that at position 935-937 holds the codon AGC rather than the codon GGC stated in SEQ ID NO: 1.

Hybridisation Protocol

The polynucleotides of the invention are such which have a nucleic acid sequence capable of hybridising with the polynucleotide sequence presented as SEQ ID NO: 1, its complementary strand, or a sub-sequence thereof, under at least medium, medium/high, or high stringency conditions, as described in more detail below.

In a preferred embodiment the polynucleotide is a fragment of at least 15 bases in length which is sufficient to permit the fragment to hybridise to DNA that encodes a polypeptide of the invention, preferably the polypeptide comprising the amino acid sequence presented as SEQ ID NO: 2 under at least medium, medium/high, or high stringency conditions, as described in more detail below.

Suitable experimental conditions for determining hybridisation between a nucleotide probe and a homologous DNA or RNA sequence, involves pre-soaking of the filter containing the DNA fragments or RNA to hybridise in 5 x SSC [Sodium chloride/Sodium citrate; cf. *Sambrook et al.*; Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Lab., Cold Spring Harbor, NY 1989] for 10 minutes, and pre-hybridisation of the filter in a solution of 5 x SSC, 5 x Denhardt's solution [cf. *Sambrook et al.*; *Op cit.*], 0.5 % SDS and 100 µg/ml of denatured sonicated salmon sperm DNA [cf. *Sambrook et al.*; *Op cit.*], followed by hybridisation in the same solution containing a concentration of 10 ng/ml of a random-primed [*Feinberg A P & Vogelstein B*; Anal. Biochem. 1983 **132** 6-13], ³²P-dCTP-labeled (specific activity > 1 x 10⁹ cpm/µg) probe for 12 hours at approximately 45°C. The filter is then washed twice for 30 minutes in 2 x SSC, 0.5 % SDS at a temperature of at least at least 60°C

(medium stringency conditions), preferably of at least 65°C (medium/high stringency conditions), more preferred of at least 70°C (high stringency conditions), and even more preferred of at least 75°C (very high stringency conditions).

Molecules to which the oligonucleotide probe hybridises under these
5 conditions may be detected using a x-ray film.

DNA Sequence Homology

In a preferred embodiment, the polynucleotides of the invention show a homology of at least 50%, preferably at least 70%, more preferred at least 80%, even
10 more preferred at least 90%, most preferred at least 95%, with the polynucleotide sequence presented as SEQ ID NO: 1.

As defined herein, the DNA sequence homology may be determined as the degree of identity between two DNA sequences indicating a derivation of the first sequence from the second. The homology may suitably be determined by means of
15 computer programs known in the art such as GAP provided in the GCG program package [*Needleman S B and Wunsch C D, Journal of Molecular Biology* 1970 **48** 443-453] using default parameters suggested herein.

Cloned Polynucleotides

20 The isolated polynucleotide of the invention may in particular be a cloned polynucleotide.

As defined herein, the term "cloned polynucleotide", refers to a polynucleotide or DNA sequence cloned in accordance with standard cloning procedures currently used in genetic engineering to relocate a segment of DNA, which
25 may in particular be cDNA, i.e. enzymatically derived from RNA, from its natural location to a different site where it will be reproduced.

Cloning may be accomplished by excision and isolation of the desired DNA segment, insertion of the piece of DNA into the vector molecule and incorporation of the recombinant vector into a cell where multiple copies or clones of the DNA
30 segment will be replicated, by reverse transcription of mRNA (reverse transcriptase technology), and by use of sequence-specific oligonucleotides and DNA polymerase in a polymerase chain reaction (PCR technology).

The cloned polynucleotide of the invention may alternatively be termed "DNA construct" or "isolated DNA sequence", and may in particular be a
35 complementary DNA (cDNA).

It is well established that potassium channels may be formed as heteromeric channels, composed of different subunits. Also it has been found that the potassium channel of the invention may form heteromers with other KCNQ's, in

particular KCNQ3, when co-expressed with these subunits. In addition, potassium channels can also associate with non-homologous subunits, e.g. the KCNE1 (formerly known as minK) subunit, that can functionally modulate these channels or lead to a specific localisation within the cell.

- 5 Therefore, in a preferred embodiment, the polynucleotide of the invention is cloned and either expressed by itself or co-expressed with polynucleotides encoding other subunits, in particular a polynucleotide encoding a KCNQ3 channel subunit.

Biological Sources

- 10 The isolated polynucleotide of the invention may be obtained from any suitable source. In a preferred embodiment, which the polynucleotide of the invention is cloned from, or produced on the basis of a cDNA library, e.g. of the retina, brain, skeletal muscle. Commercial cDNA libraries are available from e.g. Stratagene and Clontech.

- 15 The isolated polynucleotide of the invention may be obtained by methods known in the art, e.g. those described in the working examples below.

In a preferred embodiment the polynucleotide of the invention may be obtained using the PCR primers described in the working examples and presented as SEQ ID NOS: 3-32.

20

Preferred Polynucleotides

In a preferred embodiment, polynucleotide of the invention comprises the polynucleotide sequence presented as SEQ ID NO: 1.

- 25 In another preferred embodiment the polynucleotide of the invention is a sequence giving rise to KCNQ4 channels subunits comprising one or more substitutions.

In another preferred embodiment the polynucleotide of the invention is a sequence giving rise to KCNQ4 channels subunits comprising one or more substitutions in the conserved regions, as defined in more details below.

- 30 In a more preferred embodiment the polynucleotide of the invention has a polynucleotide sequence giving rise to the G285S mutation as indicated in SEQ ID NO: 1, e.g. the DNA sequence that at position 935-937 holds the codon AGC rather than the codon GGC stated in SEQ ID NO: 1.

- 35 Also contemplated within the scope of this invention are the primer sequences used in Example 2 below for the amplification of the single KCNQ4 exons, that can then be screened for mutations.

Therefore, in another preferred embodiment the polynucleotide of the invention is a primer sequence comprising any one of the polynucleotide sequences presented as SEQ ID NOS: 3-32.

It has been demonstrated that KCNQ channels often show alternative
 5 splicing and therefore may occur as isoforms originating from the same gene. Such isoforms as well as the different cDNA sequences from which they occurred are also contemplated within the scope of the present invention.

Finally the genes encoding KCNQ channel subunits in other species have been found to differ slightly from the human genes. However, genes of other species,
 10 e.g. mouse, rat, monkey, rabbit, etc., are also contemplated within the scope of the present invention.

Recombinantly Produced Polypeptides

In another aspect the invention relates to substantially pure functional
 15 polypeptides that have the electrophysiological and pharmacological properties of a KCNQ4 channel, or KCNQ4 channel subunits. The novel polypeptides of the invention may be obtained by the polynucleotides of the invention using standard recombinant DNA technology.

In a preferred embodiment, a polypeptide of the invention is the KCNQ4
 20 potassium channel subunit comprising the amino acid sequence presented as SEQ ID NO: 2, and biologically active fragments hereof.

Modifications of this primary amino acid sequence may result in proteins which have substantially equivalent activity as compared to the unmodified counterpart polypeptide, and thus may be considered functional analogous of the
 25 parent proteins. Such modifications may be deliberate, e.g. as by site-directed mutagenesis, or they may occur spontaneous, and include splice variants, isoforms, homologues from other species, and polymorphisms, and include the variant KCNQ4/G285S, that is described in more detail below. Such functional analogous are also contemplated according to the invention.

Moreover, modifications of this primary amino acid sequence may result in
 30 proteins which do not retain the biological activity of the parent protein, including dominant negative forms, etc. A dominant negative protein may interfere with the wild-type protein by binding to, or otherwise sequestering regulating agents, such as upstream or downstream components, that normally interact functionally with the
 35 polypeptide. Such dominant negative forms are also contemplated according to the invention.

In the context of this invention, the term "variant polypeptide" means a polypeptide (or protein) having an amino acid sequence that differs from the sequence

presented as SEQ ID NO: 2 at one or more amino acid positions. Such variant polypeptides include the modified polypeptides described above, as well as conservative substitutions, splice variants, isoforms, homologues from other species, and polymorphisms, and includes the variant KCNQ4/G285S (i.e. KCNQ4/G333S according to the KCNQ1 numbering).

As defined herein, the term "conservative substitutions" denotes the replacement of an amino acid residue by another, biologically similar residue. Examples of conservative substitutions include the substitution of one hydrophobic residue such as isoleucine, valine, leucine or methionine for another, or the substitution of one polar residue for another, such as the substitution of arginine for lysine, glutamic for aspartic acid, or glutamine for asparagine, and the like. The term conservative substitution also include the use of a substituted amino acid residue in place of an un-substituted parent amino acid residue provided that antibodies raised to the substituted polypeptide also immunoreact with the un-substituted polypeptide.

Also contemplated within the scope of this invention are the oligopeptides encoded by the primer sequences used in Example 2 below for the amplification of the single KCNQ4 exons, that can then be screened for mutations.

KCNQ1 Numbering System

In the context of this invention, amino acid residues (as well as nucleic acid bases) are specified using the established one-letter symbol.

By aligning the amino acid sequences of a polypeptide of the present invention to those of the known polypeptides, a specific amino acid numbering system may be employed, by which system it is possible to unambiguously allot an amino acid position number to any amino acid residue in any KNCQ channel protein, which amino acid sequence is known.

Such an alignment is presented in Table 1, below. Using the ClustalX computer alignment program [Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, & Higgins DG: The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools; Nucleic Acids Res. 1997 **25** (24) 4876-82], and the default parameters suggested herein, the amino acid sequence of a polypeptide of the present invention (hKCNQ4) and the amino acid sequences of the known polypeptides hKCNQ2-3 are aligned with, and relative to, the amino acid sequences of the known polypeptide hKCNQ1 (formerly known as KvLQT1). In the context of this invention this numbering system is designated the KCNQ1 Numbering System.

In describing the various protein variants produced or contemplated according to the invention, the following nomenclatures have been adapted for ease of reference:

Original amino acid / Position / Substituted amino acid

According to this nomenclature the substitution of serine for glycine at position 333 is designated as "G333S".

5 A deletion of glycine at the same position is designated "G333*".

An insertion of an additional amino acid residue, in this example lysine, may be designated "G333GK" or "*334K" (assumed that no position exists for this position in the amino acid sequence used for establishing the numbering system).

10 An insertion of an amino acid residue, in this example valine, at a position which exists in the established numbering system, but where no amino acid residue is actually present, may be designated "-301V".

Table 1
CLUSTAL X Multiple Sequence Alignment

15 **KCNQ1 Numbering**

hKCNQ2	MVQKSR----	-----	NGGVYPGPSG	EKKLKVG---	-FVGLDPG--	-----	APD	60
hKCNQ3	MGLKARRAAG	AAGGGGDGGG	GGGGAANPAG	GDAAGDEE	RKVGLAPGDV	EQVTLALGAG		
hKCNQ4	MAEAPPR---	-----	RLGLGPPPGD	APRAELVALT	-AVQSEQGE-	-----	AGG	
20 hKCNQ1	MAAASSPPR-	-AE--RKR-W	GWGRLPGARR	GSAGLAKKCP	FSLELAEG--	-----	GPA	
	*	.	*	.	.	:	*	.
hKCNQ2	STRDGALLIA	G-----S	EAPKRGSI	LKPRAGGAGAG	KPPKRN-AFY	RK-----	L	120
hKCNQ3	ADKDGTLLLE	GGG-----R	DEGQRRTPQG	IGLLAKTPLS	RPVKRNNAKY	RR-----	I	
25 hKCNQ4	GGSPRRLGLL	G-----S	PLPPGAPLPG	PGSGSGSACG	QRSSAAHKRY	RR-----	L	
hKCNQ1	GGALYAPIAP	GAPGPAPPAS	PAAPAAPPVA	SDLGPRPPVS	LDPRVSIYST	RRPVLARTHV		
	.	*	.	.	.	:	*	:
hKCNQ2	QNFLYNVLER	PRGW-AFIYH	AYVFLLVFSC	LVLSVVFSTIK	EYEKSSEGAL	YILEIVTIVV		180
30 hKCNQ3	QTLIYDALER	PRGW-ALLYH	ALVFLIVLGC	LILAVLTTFK	EYETVSGDWL	LLLETFAIFI		
hKCNQ4	QNWVYNVLER	PRGW-AFVYH	VFIFLLVVFSC	LVLSVLSTIQ	EHQELANECL	LILEFVMIVV		
hKCNQ1	QGRVYNFLER	PTGWKCFVYH	FAVFLIVLVC	LIFSVLSTIE	QYAALATGTL	FWMEIVLVVF		
	*	:*	***	* **	..:***	:**.*:	*	:* :...:
35 hKCNQ2	FGVEYFVRIW	AAGCCCRYRG	WRGRLKFARK	PFCVIDIMVL	IASIAVLAAG	SQGNVFATSA		240
hKCNQ3	FGAEFALRIW	AAGCCCRYKG	WRGRLKFARK	PLCMLDIFVL	IASVPVAVG	NQGNVLATS-		
hKCNQ4	FGLEYIVRVW	SAGCCCRYRG	WQGRFRFARK	PFCVIDFIVF	VASVAVIAAG	TQGNIFATSA		
hKCNQ1	FGTEYVRLW	SAGCRSKYVG	LWGRLRFARK	PISIIDLIVV	VASMVLCVG	SKGQVFATSA		
	** *	:*:*	:***	.* *	**:*:****	*:..:***:	:***	:*..* :*:::***

hKCNQ2 LRSLRFLQIL RMIRMDRRGG TWKLLGSVVY AHSKELVTAW YIGFLCLILA SFLVYLAEK- 300
hKCNQ3 LRSLRFLQIL RMLRMDRRGG TWKLLGSAIC AHSKELITAW YIGFLTLILS SFLVYLVEKD
hKCNQ4 LRSMRFLQIL RMVRMDRRGG TWKLLGSVVY AHSKELITAW YIGFLVLIFA SFLVYLAEKD
hKCNQ1 IRGIRFLQIL RMLHVDRQGG TWRLGSVVF IHRQELITTL YIGFLGLIFS SYFVYLAEKD
5 :*.:***** **::**:* **::*****: * :*:*: ***** **:: *:***.**

hKCNQ2 -----GE- -NDHFDTYAD ALWWGLITLT TIGYGDKYPQ TWNGRLLAAT FTLIGVSFFA 360
hKCNQ3 VPEVDAQGEE MKEEFETYAD ALWWGLITLA TIGYGDKTPK TWEGRLIAAT FSLIGVSFFA
hKCNQ4 ----- ANSDFSSYAD SLWWGTITLT TIGYGDKTPH TWLGRVLAAG FALLGISFFA
10 hKCNQ1 -----AVNES GRVEFGSYAD ALWWGVVTVT TIGYGDKVPQ TWVGKTIASC FSVFAISFFA
. . * :*** :***** :*: ***** *: ** *: **: *:***:****

hKCNQ2 LPAGILGS GF ALKVQEQHRQ KHFEKRRNPA AGLIQSAWRF YATNLSRTDL HSTWQYYERT 420
hKCNQ3 LPAGILGS GL ALKVQEQHRQ KHFEKRRKPA AELIQAAWRY YATNPNRIDL VATWRFYESV
15 **hKCNQ4 LPAGILGS GF ALKVQEQHRQ KHFEKRRMPA ANLIQAAWRL YSTDMSRAYL TATWYYYDSI**
hKCNQ1 LPAGILGS GF ALKVQQKQORQ KHFNQRIPAA ASLIQTAWRC YAAE--NPD- SSTWKIYIRK
*****: *****::** ***::: . * * ***:*** *:*** . :** *

hKCNQ2 VT----- VPMYRLIPP- -LNQLELLRN LKSKSGLAFR K----- DPPPEPSPSQ 480
20 hKCNQ3 VS----- FPFRRKE--- ---QLEAAS- -----S--Q
hKCNQ4 LPSFRELALL FEHVQRARNG GLRPLEVRRR PVPDGAPSRV PPVATCHRPG STSFCPGESS
hKCNQ1 AP----- RSHTLLS--- ---PSPKPKK -----S
.

hKCNQ2 KVSLKDRV-F SSPRGVAAKG KGSPQAQTVR RSPSADQSLE D-SPSKVPKSW SFG-DRSRA 540
25 hKCNQ3 KLGLLDRVRL SNPRGSNTKG K-----L FTPLNVDAIE E-SPSKEPKPV GLN-NKERF
hKCNQ4 RMGIKDRIRM GSSQRRTGPS KQQLAPPTMP TSPSSEQVGE ATSPTKVQKSW SFN-DRTRF
hKCNQ1 VVVKKKKFKL DKDNGVTPGE K-MLTVPHIT CDPPEERRLD HFSVDGYDSSV RKSPTLLEV
: . . . : . . * * : * . . .

hKCNQ2 RQAFRIKGAA SRQNSEEASL PGEDIVDDKS CPCEFVTEDL TPGLKVSIRA VCVMRFLVSK 600
hKCNQ3 RTAFRMKAYA FWQSSSEDAGT -GDPMEDRG YGNDFFIEDM IPTLKAIRA VRILQFRLYK
hKCNQ4 RASLRLLK-- -RTSAEDAPS --EEVAEEKS YQCELTVDI MPAVKTVIRS IRILKFLVAK
hKCNQ1 SMPHFMRT-- -NSFAEDLDL EGETLLTPIT H-----ISQL REHHRATIKV IRRMQYFVAK
. :: :*: : : .:: :. *: : :*: *

35 hKCNQ2 RKFKESLRPY DVMDVIEQYS AGHLDMLSRI KSLQSRVDQI VGRGP----A ITDKDR-TK 660
hKCNQ3 KKFKETLRPY DVKDVIEQYS AGHLDMLSRI KYLQTRIDMI FTPGP----P STPKHKKSQK
hKCNQ4 RKFKETLRPY DVKDVIEQYS AGHLDMLGRI KSLQTRVDQI VGRGP----G DRKAREKGDK
hKCNQ1 KKFKQARKPY DVRDVIEQYS QGHLNLMVRI KELQRRILDQS IGK-PSLFIS VSEKSK--DR
40 **:*** :** ** ***** ***::: ** * ** *: * . * . :

hKCNQ2	G-----	-----	PAAELPEDP	SMMGRLGKVE	KQVLSMEKKL	DFLVNIY--M	720
hKCNQ3	GSAFTFPSQQ	SPRNEPYVAR	PSTSEI-EDQ	SMMGKFKVKE	RQVQDMGKKL	DFLVDMH--M	
hKCNQ4	G-----	-----	PSDAEVVDEI	SMMGRVVKVE	KQVQSIEHKL	DLLLGFYSRC	
5 hKCNQ1	G-----	-----	-----SN	TIGARLNRVE	DKVTQLDQRL	ALITDML---	
	*		.	::	::	::	
hKCNQ2	QRMGIPPTET	EAYFGAKEPE	PAPPYHSPED	SREHVDRHGC	IVKIVRSSSS	TGQKNF---S	780
hKCNQ3	QHMERLQVQV	TEYYPTKGTS	SPAEAEKKED	NR-YSDLKTI	ICNYSETGPP	EPPYSFHQVT	
10 hKCNQ4	LRSGTSASLG	AVQVPLFDPD	ITSDYHSPVD	HE-DISVSAQ	TLISIRSVST	NMD-----	
hKCNQ1	-HQLLSLHGG	STP-GSGGPP	REGGAHITQP	CGSGGSVDPE	LFLPSNTLPT	YEQLTV----	
	:	::	
hKCNQ2	APPAAPPVQC	PPSTSWQPQS	HPRQG--HGT	SPVGDHGSLV	RIPPPPAHER	SLSAYGGGNR	840
15 hKCNQ3	IDKVSPYGFF	AHDPVNLPRG	GPSSGKVQAT	PPSSATTYVE	RPTVLPILTL	LDSRVSCHSQ	
hKCNQ4	-----	-----	-----	-----	-----	-----	
hKCNQ1	PRRGPDEGS-	-----	-----	-----	-----	-----	
20 hKCNQ2	ASMEFLRQED	TPGCRPPEGT	LRSDTSISI	PSVDHEELER	SFSGFSISQS	KENLDALNSC	900
hKCNQ3	ADLQGP-YSD	RISPRQRRSI	TRSDTPLSL	MSVNHEELER	SPSGFSISQD	RD--DYVFGP	
hKCNQ4	-----	-----	-----	-----	-----	-----	
hKCNQ1	-----	-----	-----	-----	-----	-----	
25							
hKCNQ2	YAAVAPCAKV	RPYIAEGESD	TSDSLCTPCG	PPRSATGEG	PFGDVGWAGP	RK--	954
hKCNQ3	NGGSSWMREK	R-YLAGETD	TDTPFTPSG	SMPLSSTGDG	-ISDSVWTPS	NKPI	
hKCNQ4	-----	-----	-----	-----	-----	-----	
hKCNQ1	-----	-----	-----	-----	-----	-----	

30

35

40

hKCNQ1: Human KCNQ1 [Wang, Q *et al.*, Nature Genet. 1996 **12** 17-23]

hKCNQ2: Human KCNQ2 [Biervert *et al.*, Science 1998 **279** 403-406]

hKCNQ3: Human KCNQ3 [Schroeder *et al.*, Nature 1998 **396** 687-690]

hKCNQ4: Human KCNQ4; A protein of the invention

- No amino acid in this position.

* Indicates positions which have a single, fully conserved residue (Conserved regions).

Biological Activity

The polynucleotide of the invention encodes a potassium channel subunit, which has been termed KCNQ4. In the cochlea, it is differentially expressed in sensory outer hair cells. A mutation in this gene in a pedigree with autosomal dominant hearing loss changes a residue in the KCNQ4 pore region. It abolishes the outwardly rectifying potassium currents of wild-type KCNQ4 on which it exerts a strong dominant negative effect.

Ion channels are excellent targets for drugs. KCNQ4, or heteromeric channels containing the KCNQ4 subunit, may be a particularly interesting target for the treatment of tinnitus and the prevention or treatment of progressive hearing loss.

KCNQ Channels in Genetic Disease

It is remarkable that mutations in every known *KCNQ* gene lead to human disease: Mutations in *KCNQ1* (KvLQT1) cause the autosomal dominant long QT syndrome (LQTS), and, when present on both alleles, the Jervell and Lange-Nielsen (JLN) syndrome whose symptoms include deafness in addition to cardiac arrhythmias. Mutations in either *KCNQ2* or *KCNQ3*, which form heteromers that probably represent the M-channel, cause benign familial neonatal convulsions (BFNC). The present invention adds *KCNQ4* and the associated autosomal dominant deafness to that list.

After *KCNQ1*, *KCNQ4* is now the second *KCNQ* channel whose loss of function leads to deafness.

Therefore, in a preferred embodiment of the invention, mutated polynucleotides may be employed in the screening for drugs that affect diseases associated with such mutations in the *KCNQ4* gene.

Heteromers Formed by KCNQ Subunits

The *KCNQ* channels described so far function physiologically as heteromers. *KCNQ1* associates with *KCNE1* (formerly known as minK), and *KCNQ2* and *KCNQ3* form heteromeric channels that underlie the M-current, an important determinant of neuronal excitability that is regulated by several neurotransmitters.

Like other *KCNQ* channel subunits, *KCNQ4* may interact with other subunits, e.g. *KCNE1* or other *KCNQ* channel subunits, and in particular with *KCNQ3*. Currents from homomeric *KCNQ3* are very small and often cannot be distinguished from *Xenopus* oocyte background currents. Co-expression of *KCNQ3* with *KCNQ4* markedly increased current amplitudes. Significantly, heteromeric *KCNQ3/KCNQ4* channels activated faster than homomeric *KCNQ4* channels, the voltage-dependence was shifted to more negative potentials, and currents displayed a different drug sensitivity.

Antibodies

The polypeptides of the invention can be used to produce antibodies which are immunoreactive or bind to epitopes of these polypeptides. Antibodies which consist essentially of pooled monoclonal antibodies with different specificities, as well as distinct monoclonal antibody preparations may be provided.

The preparation of polyclonal and monoclonal antibodies is well known in the art. Polyclonal antibodies may in particular be obtained as described by e.g. *Green et al.*: "Production of Polyclonal Antisera" in Immunochemical Protocols (Manson, Ed.); Humana Press, 1992, Pages 1-5; and *Coligan et al.*: "Production of Polyclonal Antisera in rabbits, rats, Mice and Hamsters" in Current Protocols in Immunology, 1992, Section 2.4.1; which protocols are hereby incorporated by reference.

Monoclonal antibodies may in particular be obtained as described by e.g. *Kohler & Milstein*, Nature 1975 **256** 495; *Coligan et al.* in Current Protocols in Immunology, 1992, Sections 2.5.1 - 2.6.7; and *Harlow et al.* in Antibodies: A Laboratory Manual; Cold Spring Harbor Pub., 1988, Page 726; which protocols are hereby incorporated by reference.

Briefly, monoclonal antibodies may be obtained by injecting e.g. mice with a composition comprising an antigen, verifying the presence of antibody production by removing a serum sample, removing the spleen to obtain B lymphocytes, fusing the B lymphocytes with myeloma cells to produce hybridomas, cloning the hybridomas, selecting positive clones that produce the antibodies to the antigen, and isolating the antibodies from the hybridoma cultures.

Monoclonal antibodies can be isolated and purified from hybridoma cultures by a variety of well-established techniques, including affinity chromatography with protein A Sepharose, size-exclusion chromatography, and ion-exchange chromatography, see. e.g. *Coligan et al.* in Current Protocols in Immunology, 1992, Sections 2.7.1 - 2.7.12, and Sections 2.9.1 - 2.9.3; and *Barnes et al.*: "Purification of Immunoglobulin G (IgG)" in Methods in Molecular Biology; Humana Press, 1992, Vol. 10, Pages 79-104.

The polyclonal or monoclonal antibodies may optionally be further purified, e.g. by binding to and elution from a matrix to which the polypeptide, to which the antibodies were raised, is bound.

Antibodies which bind to the polypeptide of the invention can be prepared using an intact polypeptide or fragments containing small peptides of interest as the immunising antigen. The polypeptide used to immunise an animal may be obtained by recombinant DNA techniques or by chemical synthesis, and may optionally be conjugated to a carrier protein. Commonly used carrier proteins which are chemically

coupled to the peptide include keyhole limpet hemocyanin (KLH), thyroglobulin, bovine serum albumin (BSA), and tetanus toxoid. The coupled peptide may then be used to immunise the animal, which may in particular be a mouse, a rat, a hamster or a rabbit.

5

Genetically Manipulated Cells

In a third aspect the invention provides a cell genetically manipulated by the incorporation of the heterologous polynucleotide of the invention. The cell of the invention may in particular be genetically manipulated to transiently or stably express,
10 over-express or co-express a KCNQ4 channel subunit as defined above. Methods of transient and stable transfer are known in the art.

The polynucleotide of the invention may be inserted into an expression vector, e.g. a plasmid, virus or other expression vehicle, and operatively linked to expression control sequences by ligation in a way that expression of the coding
15 sequence is achieved under conditions compatible with the expression control sequences. Suitable expression control sequences include promoters, enhancers, transcription terminators, start codons, splicing signals for introns, and stop codons, all maintained in the correct reading frame of the polynucleotide of the invention so as to permit proper translation of mRNA. Expression control sequences may also include
20 additional components such as leader sequences and fusion partner sequences.

The promoter may in particular be a constitutive or an inducible promoter. When cloning in bacterial systems, inducible promoters such as pL of bacteriophage γ , plac, ptrp, ptac (ptrp-lac hybrid promoter), may be used. When cloning in mammalian systems, promoters derived from the genome of mammalian cells, e.g.
25 the TK promoter or the metallothionein promoter, or from mammalian viruses, e.g. the retrovirus long terminal repeat, the adenovirus late promoter or the vaccinia virus 7.5K promoter, may be used. Promoters obtained by recombinant DNA or synthetic techniques may also be used to provide for transcription of the polynucleotide of the invention.

30 Suitable expression vectors typically comprise an origin of expression, a promoter as well as specific genes which allow for phenotypic selection of the transformed cells, and include vectors like the T7-based expression vector for expression in bacteria [*Rosenberg et al*; Gene 1987 **56** 125], the pMSXND expression vector for expression in mammalian cells [*Lee and Nathans*, J. Biol. Chem. 1988 **263**
35 3521], baculovirus derived vectors for expression in insect cells, and the oocyte expression vector PTLN [*Lorenz C, Pusch M & Jentsch T J*: Heteromultimeric CLC chloride channels with novel properties; Proc. Natl. Acad. Sci. USA 1996 **93** 13362-13366].

In a preferred embodiment, the cell of the invention is an eukaryotic cell, in particular a mammalian cell, an oocyte, or a yeast cell. In a more preferred embodiment, the cell of the invention is a human embryonic kidney (HEK) cell, a HEK 293 cell, a BHK21 cell, a Chinese hamster ovary (CHO) cell, a *Xenopus laevis* oocyte
 5 (XLO) cell, a COS cell, or any other cell line able to express KCNQ potassium channels.

When the cell of the invention is an eukaryotic cell, incorporation of the heterologous polynucleotide of the invention may be in particular be carried out by infection (employing a virus vector), by transfection (employing a plasmid vector), or
 10 by calcium phosphate precipitation, microinjection, electroporation, lipofection, or other physical-chemical methods known in the art.

In a further preferred embodiment, the cell of the invention is genetically manipulated to co-express KCNQ4 and KCNQ1 channel subunits; KCNQ4 and KCNQ2 channel subunits; KCNQ4 and KCNQ3 channel subunits; KCNQ4 and
 15 KCNQ1 and KCNQ2 channel subunits; KCNQ4 and KCNQ1 and KCNQ3 channel subunits; KCNQ4 and KCNQ2 and KCNQ3 channel subunits; or KCNQ4 and KCNQ1 and KCNQ2 and KCNQ3 channel subunits.

KCNQ4 Active Chemical Compounds

20 In another aspect the invention relates to chemical compounds capable of binding to, and showing activity at potassium channels containing one or more KCNQ4 subunits. In the context of this invention such compounds are termed KCNQ4 active compounds. The KCNQ4 active compounds of the invention show activity in concentrations below 100 μ M, preferably below 10 μ M, more preferred below 1 μ M. In
 25 its most preferred embodiment the KCNQ4 active compounds of the invention show activity in low micromolar and the nanomolar range.

The KCNQ4 active compounds of the invention have therapeutic potential, and may be used for the manufacture of pharmaceutical compositions.

The KCNQ4 active compounds of the invention may in particular be used in
 30 diagnosis, treatment, prevention or alleviation of diseases related to tinnitus, loss of hearing, in particular progressive hearing loss, neonatal deafness, and presbycusis (deafness of the elderly); and diseases or adverse conditions of the CNS, including affective disorders, Alzheimer's disease, anxiety, ataxia, CNS damage caused by trauma, stroke or neurodegenerative illness, cognitive deficits, compulsive behaviour,
 35 dementia, depression, Huntington's disease, mania, memory impairment, memory disorders, memory dysfunction, motion disorders, motor disorders, neurodegenerative diseases, Parkinson's disease and Parkinson-like motor disorders, phobias, Pick's disease, psychosis, schizophrenia, spinal cord damage, stroke, and tremor.

Currently two compound have been identified. As a preferred embodiment the invention therefore provides 1,3-dihydro-1-phenyl-3,3-bis(4-pyridylmethyl)-2H-indol-2-one (Linopirdine) and 10,10-bis(4-pyridinyl-methyl)-9(10H)-anthracenone (XE991) for use in the manufacture of a pharmaceutical composition for the diagnosis, treatment, prevention or alleviation of the above diseases.

Screening of Drugs

In a further aspect the invention provides methods for screening for KCNQ4 active compounds, i.e. chemical compounds capable of binding to, and showing activity at potassium channels containing one or more KCNQ4 subunits. The activity determined may be inhibitory activity, stimulating activity, or other modulatory activity. In particular the KCNQ4 active compound may induce a second messenger response, which cause a change of the molecular characteristics of the cell, e.g. the ion flux, enzyme activation, changes in cyclic nucleotides such as cAMP, cADP, cGMP, and cGDP, etc.

Therefore, in another aspect, the invention provides a method for identifying functional ligands for a human potassium channel, comprising a KCNQ4 subunit, which method comprises transfecting cells with one or more polypeptides of the invention, encoding a KCNQ4 channel subunit, and detecting the effect on the signal transduction pathway caused in these cells by binding of the ligands to the receptor by a reporter system.

Such chemical compounds can be identified by one of, or both methods described below.

Binding Studies

Binding studies are usually carried out by subjecting the target to binding with a labelled, selective agonist (binding agent), to form a labelled complex, followed by determination of the degree of displacement caused by the test compound upon addition to the complex.

In a specific aspect the invention provides a method of screening a chemical compound for capability of binding to a potassium channel comprising at least one KCNQ4 channel subunit, which method comprises the steps of (i) subjecting a KCNQ4 channel subunit containing cell to the action of a KCNQ4 binding agent to form a complex with the KCNQ4 channel subunit containing cell; (ii) subjecting the complex of step (i) to the action of the chemical compound to be tested; and (iii) detecting the displacement of the KCNQ4 binding agent from the complex with the KCNQ4 channel subunit containing cell.

The KCNQ4 channel subunit containing cell preferably is a cell of the invention as described above.

The KCNQ4 binding agent preferably is a radioactively labelled 1,3-dihydro-1-phenyl-3,3-bis(4-pyridylmethyl)-2H-indol-2-one (Linopirdine); or 10,10-bis(4-pyridinyl-methyl)-9(10H)-anthracenone.

In a even more preferred embodiment, the binding agent is labelled with ^3H , and the displacement of the KCNQ4 binding agent from the complex with the KCNQ4 channel subunit containing cell is detected by measuring the amount of radioactivity by conventional liquid scintillation counting.

Activity Studies

The KCNQ4 channel agonists may affect the potassium channel in various ways. The agonist may in particular show inhibitory activity, stimulating activity, or other modulatory activity.

In a specific aspect the invention provides a method for determining the activity at potassium channels containing one or more KCNQ4 subunits. According to this method a KCNQ4 channel subunit containing cell is subjected to the action of the chemical compound to be tested, and the activity is detected by way of monitoring the membrane potential, the current, the potassium flux, or the secondary calcium influx of the KCNQ4 channel subunit containing cell, preferably a genetically manipulated as described above.

The membrane potential and the current may be monitored by electrophysiologic methods, including patch clamp techniques, such as current clamp technology and two-electrode voltage clamp technology, or by spectroscopic methods, such as fluorescence methods.

In a preferred embodiment, monitoring of the membrane potential of the KCNQ4 channel subunit containing cell is performed by patch clamp techniques.

In another preferred embodiment, monitoring of the membrane potential of the KCNQ4 channel subunit containing cell is performed by spectroscopic methods, e.g. using fluorescence methods. In a more specific embodiment, the KCNQ4 channel subunit containing cell is mixed with a membrane potential indicating agent, that allow for a determination of changes in the membrane potential of the cell, caused by the addition of the test compound. The membrane potential indicating agent may in particular be a fluorescent indicator, preferably DIBAC₄(3), DiOC₅(3), and DiOC₂(3).

In yet a preferred embodiment, monitoring of the membrane potential of the KCNQ4 channel subunit containing cell is performed by spectroscopic methods, e.g. using a FLIPR assay (Fluorescence Image Plate Reader; available from Molecular Devices).

Screening of Genetic Material

In a further aspect the invention relates to the use of a polynucleotide sequence of the invention for the screening of genetic materials. By this method, 5 individuals bearing a gene identical or homologous to a polynucleotide of the invention may be identified.

In the screening method of the invention, a polynucleotide of the invention, or any fragment or sub-sequence hereof, and in particular any one of the polynucleotide sequences presented as SEQ ID NOS: 3-32, is employed. For the 10 identification of individuals bearing mutated genes, preferably a mutated form of the polynucleotide represented by SEQ ID NO: 1 is employed, and in particular a polynucleotide sequence holding the mutation giving rise to the KCNQ4/G285S variant.

In the screening method of the invention only short sequences needs to be 15 employed depending on the actual method used. For SSCA, several hundreds of base pairs may be needed, for oligonucleotide or PCR hybridisation only of from about 10 to about 50 basepairs may be needed.

In a more specific embodiment, the primer sequences used in Example 2 below for the amplification of the single KCNQ4 exons, and presented as SEQ ID 20 NOS: 3-32, may be used for the screening of mutations.

The screening may be accomplished by conventional methods, including hybridisation, SSCA analysis, and array technology (DNA chip technology). The hybridisation protocol described above represents a suitable protocol for use in a screening method of the invention.

25 Therefore, in particular embodiment the invention provides a method for the identification, localisation, isolation or amplification a polynucleotide of the invention, which method a polynucleotide primer of the invention, in particular any one of those presented as SEQ ID NOS: 3-32, is used as a probe. This method may be accomplished using conventional molecular biological techniques, e.g. those used 30 and described in the working examples below.

In a preferred embodiment, the method is used for performing gene amplification using conventional PCR techniques, e.g. as described in the working examples below.

35 Transgenic Animals

Transgenic animal models provide the means, *in vivo*, to screen for therapeutic compounds. The establishment of transgenic animals may in particular be helpful for the screening of drugs to fully elucidate the pathophysiology of

KCNQ4/DFNA2 deafness. These animals may also be valuable as a model for the frequent condition of presbycusis that also develops slowly over decades. Since *KCNQ4* is expressed also in brain, they may also be helpful in screening for drugs effective in CNS disorders, e.g. epilepsy.

5 By transgene is meant any piece of polynucleotide which is inserted by artifice into a cell, and thus becomes part of the genome of the organism that develops from that cell. Such a transgene may include a gene which is partly or entirely heterologous (i.e. foreign) to the transgenic organism, or it may represent a gene homologous to an endogenous gene of the organism.

10 By a transgenic animal is meant any organism holding a cell which includes a polynucleotide sequence which is inserted into that cell by artifice, and which cell becomes part of the transgenic organism which develops from that cell. Such a transgene may be partly or entirely heterologous to the transgenic animal. Although transgenic mice represent a preferred embodiment of the invention, other transgenic
15 mammals including, but not limited to transgenic rodents (e.g. hamsters, guinea pigs, rabbits and rats), and transgenic pigs, cattle, sheep and goats may be created by standard techniques and are included in the invention.

Preferably, the transgene is inserted by artifice into the nuclear genome.

20 Knock-out and Knock-in Animals

The transgenic knock-out animal models may be developed by homologous recombination of embryonic stem cells with constructs containing genomic sequence from the *KCNQ4* gene, that lead to a loss of function of the gene after insertion into the endogenous gene.

25 By knock-out mutation is meant an alteration in the polynucleotide sequence that reduces the biological activity of the polypeptide normally encoded therefrom. In order to create a true knock-out model, the biological activity of the expressed polypeptide should be reduced by at least 80% relative to the un-mutated gene. The mutation may in particular be a substitution, an insertion, a deletion, a
30 frameshift mutation, or a mis-sense mutation. Preferably the mutation is a substitution, an insertion or a deletion.

To further assess the role of *KCNQ4* at an organism level, the generation of an animal, preferably a mouse, lacking the intact *KCNQ4* gene, or bearing a mutated *KCNQ4* gene, is desired.

35 A replacement-type targeting vector, which may be used to create a knock-out model, may be constructed using an isogenic genomic clone, e.g. from a mouse strain such as 129/Sv (Stratagene Inc., La Jolla, CA). The targeting vector may be introduced into a suitably-derived line of embryonic stem (ES) cells by electroporation

to generate ES cell lines that carry a profoundly truncated form of the *KCNQ4* gene. The targeted cell lines may then be injected into a mouse blastula stage embryo to generate chimeric founder mice. Heterozygous offspring may be interbred to homozygosity.

As the slowly progressive hearing loss observed in DFNA2 may require the expression from one allele of a dominant negative mutant, it may also be desired to create a knock-in animal in which the wild-type *KCNQ4* gene is replaced by this mutated gene.

Animal models for over-expression may be generated by integrating one or more polynucleotide sequence of the invention into the genome according to standard techniques.

The procedures disclosed herein involving the molecular manipulation of nucleic acids are known to those skilled in the art, and are described by e.g. *Fredrick MA et al.* [*Fredrick MA et al.*: Short Protocols in Molecular Biology; John Wiley and Sons, 1995] and *Sambrook et al.* [*Sambrook et al.*: Molecular Cloning: A Laboratory Manual; 2. Ed., Cold Spring Harbor Lab.; Cold Spring Harbor, NY 1989], and in *Alexandra LJ* (Ed.): Gene Targeting: A practical approach; Oxford University Press (Oxford, New York, Tokyo), 1993.

BRIEF DESCRIPTION OF THE DRAWINGS

The present invention is further illustrated by reference to the accompanying drawing, in which:

Fig. 1 shows the electrophysiological properties of *KCNQ4* currents:

Fig. 1A: Two-electrode voltage-clamp current traces from a *Xenopus* oocyte injected with *KCNQ4* cRNA. Starting from a holding potential of -60 mV cells were clamped for 4 seconds to voltages between $-80...+60$ mV in $+10$ mV steps, followed by a constant test pulse to -30 mV;

Fig. 1B: Current traces showing the inactivation behaviour of *KCNQ4* at different voltages. After an activating voltage pulse at $+40$ mV of 3.5 seconds duration the cell was clamped to voltages between $+40...-120$ mV in -10 mV steps;

Fig. 1C: Apparent open-probability (p_{open}) as a function of voltage determined from tail current analysis of currents as in (A). Half-maximal p_{open} is archived at (-10.0 ± 1.2) mV, and the apparent gating charge is 1.4 ± 0.1 , as obtained from a fit of a Boltzmann-function to the data ($n=14$ from 2 oocyte batches, \pm S.E.M.);

Fig. 1D: Shift of the reversal potential with the extracellular K^+ -concentration with ND98 as reference solution. Total mono-valent cation concentration was 100 mM and the stated K^+ -concentration was obtained by mixing solutions ND100 and KD100. The reversal potential shift of 46.7 ± 0.9 mV per decade indicates a channel selective for K^+ ($n=18$ from 3 oocyte batches, \pm S.E.M.). Substitution of external K^+ with other cations yielded the following permeability ratios: $P_K/P_{Na} = 52.3 \pm 4.4$, $P_K/P_{Cs} = 7.8 \pm 0.7$, and $P_K/P_{Rb} = 0.94 \pm 0.03$ (permeability sequence: $Rb^+ \sim K^+ > Cs^+ \gg Na^+$, $n=15$ from 3 oocyte batches, \pm S.E.M.);

Fig. 1E: Current traces of WT KCNQ4 (thick solid line), a 1:1 co-injection of WT KCNQ4 and KCNQ4_{G285S} mutant (thin solid line) and KCNQ4_{G285S} mutant (dotted line). KCNQ4_{G285S} currents were indistinguishable from water-injected control oocytes. From a holding potential at -60 mV the cells were voltage-clamped for 6 seconds at $+40$ mV, followed by a -30 mV step;

Fig. 1F: Mean currents, measured after clamping oocytes for 4 seconds at $+40$ mV, averaged from several experiments as in (1E) ($n=20 \dots 35$, 4 oocyte batches, \pm S.E.M.); and

Fig. 2A shows KCNQ4 co-expressed with KCNQ1;

Fig. 2B shows KCNQ4 co-expressed with KCNQ2;

Fig. 2C shows KCNQ4 co-expressed with KCNQ3;

Fig. 2D shows derived dominant negative mutants (KCNQ1_{G219S}, KCNQ2_{G279S}, KCNQ3_{G318S}) ($n=10 \dots 31$, 3 oocyte batches, \pm S.E.) Representative currents from experiments as in (C) showing altered activation kinetics for the co-injection of KCNQ4 with KCNQ3 or KCNQ3_{G318S}, respectively. From a holding potential at -60 mV the voltage was clamped for 4 s at $+40$ mV, followed by a step to -30 mV. Time constants and amplitudes obtained from two-exponential fits were: KCNQ4: $t_1=360$ ms, $A_1=-4.9$ μ A, $t_2=1700$ ms, $A_2=-0.34$ μ A; KCNQ3+KCNQ4: $t_1=120$ ms, $A_1=-6.3$ μ A, $t_2=560$ ms, $A_2=-1.3$ μ A;

Fig. 2E: Apparent p_{open} as a function of voltage for currents from oocytes co-injected with KCNQ4 and KCNQ3 cRNA, determined from tail current analysis (squares, thick solid curve). Half-maximal p_{open} is achieved at $V_{0.5} = (-19.1 \pm 2.0)$ mV, and the apparent gating charge is 1.5 ± 0.2 ($n=23$ from 3 oocyte batches, \pm S.E.M.), as obtained from a fit of a Boltzmann-function to the data. The p_{open} curve for KCNQ4 is also shown for reference (circles, thin solid curve);

Fig. 2F: Current traces recorded from an oocyte co-injected with KCNQ3 and KCNQ4 cRNA with typical M-current voltage-protocol. Starting from a holding potential at -30 mV the cell was progressively hyperpolarized for 1 s to voltages between -30 and -90 mV in -10 mV steps;

Fig. 2G: Differential effects of 200 μ M Linopirdine on KCNQ4 ($n=10$, \pm S.E.M.) and KCNQ3+KCNQ4 ($n=6$, \pm S.E.M.). Currents were measured at $+40$ mV and % current remaining with Linopirdine is shown. Upon addition of Linopirdine steady-state inhibition was reached after ~ 1 min for KCNQ4 currents, and after ~ 3 min for KCNQ3+KCNQ4 currents.

EXAMPLES

The invention is further illustrated with reference to the following examples which are not intended to be in any way limiting to the scope of the invention as claimed.

Example 1

Cloning and Characterisation of KCNQ4 cDNA

Using a KCNQ3 potassium channel partial cDNA as a probe, a human retina cDNA λ phage library (Clontech, #HL1132a) was screened, and a ≈ 1 kb cDNA encoding a protein fragment homologous to KCNQ potassium channels was isolated. It was distinct from the known members KCNQ1 (KvLQT1), KCNQ2 and KCNQ3. We named the novel gene *KCNQ4*. Overlapping cDNA's containing the entire open reading frame were obtained by re-screening the cDNA library and by extending the 5' end in RACE (rapid amplification of cDNA ends) experiments using a Marathon kit (Clontech) with human skeletal muscle cDNA. A complete cDNA was assembled and cloned into the oocyte expression vector PTLN [Lorenz C, Pusch M & Jentsch T J: Heteromultimeric CLC chloride channels with novel properties; *Proc. Natl. Acad. Sci. USA* 1996 **93** 13362-13366].

The cDNA encodes a polypeptide of 695 amino acids with a predicted mass of 77 kDa (SEQ ID NO: 2). Its overall amino-acid identity to KCNQ1, KCNQ2, and KCNQ3 is 38%, 44%, and 37%, respectively. Together with these proteins it forms a distinct branch of the superfamily of voltage-gated potassium channels. As a typical member of this gene family, KCNQ4 has 6 predicted transmembrane domains and a P-loop between transmembrane domains S5 and S6. In potassium channels, which are tetramers of identical or homologous subunits, four of these highly conserved P-loops combine to form the ion-selective pore. As other KCNQ channels, KCNQ4 has a long predicted cytoplasmic carboxy terminus that accounts for about

half of the protein. A conserved region present in the carboxy termini of KCNQ1, -2, and -3 is also present in KCNQ4 (roughly represented by exon 12).

The sequence of KCNQ4 predicts several potential sites for phosphorylation by protein kinase C. In contrast to KCNQ1 and KCNQ2, however, it lacks an amino terminal consensus site for cAMP-dependent phosphorylation.

A human multiple tissue Northern blot (Clontech, #7760-1) was probed with a 749 bp EcoRI/PmlI cDNA fragment of KCNQ4. The fragment was labelled with ³²P using the Rediprime labelling kit (Amersham). Hybridisation was performed in ExpressHyb solution according to the instructions of the manufacturer (Clontech). The filter was then exposed to Kodak BioMax film for 4 days.

Northern analysis of *KCNQ4* expression in human tissues revealed faint bands of ≈5 kb in heart, brain and skeletal muscle. In some tissues, there was also a larger band. Upon longer exposure, weaker ≈5 kb bands were also detected in other tissues including kidney and pancreas.

Example 2

Genomic structure and chromosomal mapping to the DFNA2 locus

A PAC was isolated that contains the entire *KCNQ4* coding region. The genomic structure of the *KCNQ4* gene was established (SEQ ID NO: 1).

The genomic structure was established by a PCR approach from genomic DNA. Individual *KCNQ4* exons and adjacent short intronic sequences were amplified by standard PCR techniques from human genomic DNA using intronic oligonucleotide primers.

For amplification, the following intronic primer pairs were used (all primers in 5'→3' direction; in brackets the size of the PCR product):

- | | | |
|---------------------|-------------------------|-------|
| 1a (SEQ ID NO: 3): | catgctctctgagcgccccgagc | |
| 1r (SEQ ID NO: 4): | aggccaggcttgcgcggggaacg | (544) |
| 2a (SEQ ID NO: 5): | cagcacagagctgtaactccagg | |
| 2r (SEQ ID NO: 6): | aagctgctctctgagccatgg | (500) |
| 3a (SEQ ID NO: 7): | gctgggtccgcgctgtgacc | |
| 3r (SEQ ID NO: 8): | ggtctccagggtcagagtcg | (292) |
| 4a (SEQ ID NO: 9): | tccgggtccgtgcgcggggta | |
| 4r (SEQ ID NO: 10): | gagacagcccctctgacctcg | (328) |
| 5a (SEQ ID NO: 11): | atcccttcccgtgtggaagc | |
| 5r (SEQ ID NO: 12): | agtcacgatgggcagacctcg | (286) |
| 6a (SEQ ID NO: 13): | cctcatgatcaggctcctacc | |
| 6r (SEQ ID NO: 14): | atgttgacaggggtgagc | (270) |
| 7a (SEQ ID NO: 15): | aaggatggggacacccttgc | |

	7r (SEQ ID NO: 16):	acacaggggtgacacacc	(244)
	8a (SEQ ID NO: 17):	gctctgggtaacccacaactg	
	8r (SEQ ID NO: 18):	gctccccctgggagccatcacc	(316)
	9a (SEQ ID NO: 19):	tgagctcaggagctctgtgc	
5	9r (SEQ ID NO: 20):	accacgaagtggctgaaggc	(346)
	10a (SEQ ID NO: 21):	gtcctaagtcagctttgtcc	
	10r (SEQ ID NO: 22):	cctcagccggccctcgatcg	(347)
	11a (SEQ ID NO: 23):	cactctactggtggttggc	
	11r (SEQ ID NO: 24):	ctctgacctcaagtgatcc	(281)
10	12a (SEQ ID NO: 25):	gatagcaaagagatggagagg	
	12r (SEQ ID NO: 26):	aactcagctgcagcagtgagc	(328)
	13a (SEQ ID NO: 27):	gtgccttctccttcacaggc	
	13r (SEQ ID NO: 28):	aacgcacctccccatgtca	(297)
	14a (SEQ ID NO: 29):	ttgtgcttcccagataagc	
15	14r (SEQ ID NO: 30):	cgtgagggagtgagttcaagtacg	(445)

Sequences of exons and adjacent introns are deposited in GenBank (Accession Numbers AF105203-AF105216).

To screen unlinked pedigrees with autosomal dominant deafness we amplified only exons 4 through 7 that code for the pore and adjacent transmembrane domains as these may have the highest likelihood to harbour mutations. After amplification and agarose gel purification, PCR products were directly sequenced using the amplification primers and an ABI377 automated DNA sequencer.

The highly conserved transmembrane block S1-S6 was found to be encoded by 6 exons (exons 2 to 7) having the same limits as in KCNQ2 and KCNQ3. In KCNQ1 an additional intron interrupts the sequence encoding domain S4. The exon-intron structures of KCNQ genes diverge most in the poorly conserved carboxy-termini of these proteins.

Using hybridisation to human chromosomes, *KCNQ4* was mapped to 1p34.

A PAC containing the coding sequence of *KCNQ4* was isolated using intronic *KCNQ4* oligonucleotide primers and PCR. It was used to localise *KCNQ4* to 1p34 using FISH (Genome Systems). *KCNQ4* was then mapped on the Whitehead Contig WC1.10 using several of the intronic primers given above and published STS markers by PCR amplification from individual YAC clones.

Several diseases have been mapped to this region. This includes DFNA2, a locus for dominant progressive hearing loss. Due to the critical role of K⁺ homeostasis in auditory mechano transduction, we considered *KCNQ4* as an excellent candidate gene for DFNA2. The DFNA2 locus has been mapped between markers D1S255 and D1S193. We therefore refined the localisation of *KCNQ4* in

comparison to published physical and genetic maps using a YAC (yeast artificial chromosome) contig of this region. *KCNQ4* was present on CEPH YAC clone 914c3, a result which places this gene within the DFNA2 region.

5 Example 3

Expression of *KCNQ* genes in the inner ear

The expression of *KCNQ4*, as well as of other *KCNQ* genes, was studied by semi-quantitative RT-PCR on mouse cochlear RNA.

10 RT-PCR analysis of mouse *KCNQ* mRNA expression

Approximately 2 µg of mouse total brain RNA and mouse cochlear and vestibular RNA were reverse transcribed using the SuperScript™ II (Gibco BRL) reverse transcriptase.

1 µl (resp. 1 µl of a 1:10 dilution) of cDNA was amplified for 30 cycles (96°C
15 for 30 sec, 61°C for 30 sec, and 68°C for 45 sec) using a 2400 Thermocycler System (Perkin Elmer). Each 50 µl reaction contained 2.5 U polymerase (Expand™ Long Template PCR System, Boehringer Mannheim) and 5% DMSO.

KCNQ1 primers were based on the mouse cDNA sequence (GenBank Accession # U70068):

20 MK1a 5'-aaggctggatcagtcattgg-3'; and

MK1r 5'-agggtggcaggctgttgctgg-3' (280 bp)

As no mouse *KCNQ2* sequence was available, we chose sequences conserved between human (Y15065) and rat (AF087453) *KCNQ2*:

MK2a 5'-gccacggcacctccccctgg-3'; and

25 MK2r 5'-ccctctgcaatgtaggcctgac-3' (331 bp)

KCNQ3 primers were derived from a mouse EST (AA386747):

MK3a 5'-ccaaggaatgaaccatgtagcc-3'; and

MK3r 5'-cagaagagtcaagatgggcaggac-3' (461 bp)

Mouse *KCNQ4* primers were:

30 (SEQ ID NO: 31): MK4a 5'-agtacctgatggagcgccctctcg-3'; and

(SEQ ID NO: 32): MK4r 5'-tcatccaccgtaagctcacactgg-3' (366 bp)

Amplification products were verified by direct sequencing.

These results were compared with those obtained with vestibular and brain RNA.

35 *KCNQ1*, *KCNQ3* and *KCNQ4* messages can be detected in the cochlea, and additional PCR cycles revealed a weak *KCNQ2* expression as well. At this high amplification, *KCNQ1* was also detected in brain. *KCNQ1* and *KCNQ4* appear to have the highest cochlear expression. *KCNQ1* expression is higher in the cochlea than in

brain (which was negative by Northern analysis). The reverse is true for *KCNQ2* and *KCNQ3*, both of which are broadly expressed in brain. *KCNQ4* expression is significant in both of these tissues.

5 In situ hybridisation of mouse cochlea

In situ hybridisation's were performed on cochlea sections from mice at postnatal day P12 with a *KCNQ4* antisense probe.

A mouse *KCNQ4* cDNA corresponding to bp 618 to 1602 of the human *KCNQ4* ORF was cloned into pBluescript. Sense and antisense probes were transcribed using T3 and T7 RNA polymerases after appropriate linearization. After
10 DNase digestion, the probes were ethanol precipitated twice with 0.4 M LiCl. They were labelled with digoxigenin-11-UTP as described previously (Schaeren-Wiemers and Gerfin-Moser, 1993).

Mouse inner ears were fixed for 1 hour at 4 °C in 4 % paraformaldehyde in
15 PBS. After three rinses in PBS, they were immersed in 20 % sucrose overnight at 4°C. Cryostat sections (10-14 µm) were post-fixed and rinsed in PBS. Following pre-hybridisation at room temperature for at least 3 hours, they were hybridised overnight at 58 °C in a humid chamber. Sections were then washed and incubated with sheep antidigoxigenin antibody coupled to alkaline phosphatase. Staining by NBT/BCIP
20 (Boehringer Mannheim) was done for 2 hours at 37°C and overnight at RT. Sections were then mounted in Aquatex (Merck, USA).

Sensory outer hair cells were strongly labelled. By contrast, the inner hair cells appeared negative. The stria vascularis, the site of *KCNQ1* expression, was negative as well. Control hybridisation with a *KCNQ4* sense probe revealed that the
25 staining of outer hair cells was specific.

Autosomal Dominant Deafness

These results indicated that *KCNQ4* was an excellent candidate gene for autosomal dominant deafness. As we did not have access to the published pedigrees
30 that were linked to the DFNA2 locus, we screened 45 families with autosomal dominant deafness without previous linkage analysis. In most of these families, the hearing loss had been diagnosed before adulthood, i.e. before the age of onset reported for most of the DFNA forms, including DFNA2.

Mutation screening was limited to exons 4 to 7 that encode the pore region
35 and adjacent transmembrane domains. A *KCNQ4* mutation was found in a French family with profound hearing loss. Its clinical features include progressive hearing loss that is more prominent with higher frequencies, tinnitus in one patient, and no indication for vestibular defects nor gross morphological changes in the inner ear. A

mis-sense mutation (cf. SEQ ID NO: 1; The mutation G935A at the nucleotide level giving rise to the variant G285S at the amino acid level) was present in exon 6 in a heterozygous state. Using an AluI restriction site (AGCT) introduced by this mutation, it was shown that it co-segregated with all affected members in the pedigree. This mutation was not found on 150 control Caucasian chromosomes.

The G285S mutation affects the first glycine in the GYG signature sequence of potassium channel pores. This glycine is highly conserved across different classes of potassium channels in all species. The crystal structure of the *Streptomyces lividans* potassium channel reveals that these three amino acids line the narrowest part of the ion-conductive pore. Mutations in these amino-acids disrupt the selectivity filter and in most cases lead to a loss of channel function. Interestingly, an identical change in amino acids at the equivalent position was found in the *KCNQ1* gene of a patient with the dominant long QT syndrome. It disrupted channel activity and exerted a dominant negative effect on co-expressed WT *KCNQ1* channels. These mutations also have dominant negative effects when inserted into *KCNQ2* and *KCNQ3*. This is strong evidence that the progressive hearing loss in this family is due to the *KCNQ4*/G285S mutation.

Example 4

20 Functional expression of *KCNQ4* potassium channel subunits

KCNQ4 was expressed in *Xenopus* oocytes and its activity was investigated by two-electrode voltage clamping.

After linearization of the *KCNQ4*-containing PTLN vector with HpaI, capped cRNA was transcribed *in vitro* using the mMessage mMachine cRNA synthesis kit (Ambion). Usually 5 – 15 ng of cRNA were injected into *Xenopus* oocytes previously isolated by manual defolliculation and short collagenase treatment. In co-expression experiments cRNAs were injected at a 1:1 ratio. Oocytes were kept at 17°C in modified Barth's solution (90 mM NaCl, 1 mM KCl, 0.41 mM CaCl₂, 0.33 mM Ca(NO₃)₂, 0.82 mM MgSO₄, 10 mM HEPES, 100 U penicillin–100 µg streptomycin/ml, pH 7.6).

Standard two-electrode voltage-clamp measurements were performed at room temperature 2-4 days after injection using a Turbotec 05 amplifier (npi instruments, Tamm, Germany) and pClamp 5.5 software (Axon Instruments). Currents were usually recorded in ND98 solution (see Table 2). Solutions for Na⁺ / K⁺ replacement experiments were prepared from an appropriate mixture of solution KD100 (100 mM KCl) and ND100 (100 mM NaCl) to yield the stated concentrations of Na⁺ and K⁺. Linopirdine (RBI, Natick, MA) was prepared as a 100 mM stock solution in DMSO and added to a final concentration of 200 µM to ND98.

Table 2
Solution contents (concentrations in mM)

ND98	ND 100	KD100	Rb100	Cs100
98 NaCl 2 KCl 0.2 CaCl ₂ 2.8 MgCl ₂	100 NaCl 0.2 CaCl ₂ 2.8 MgCl ₂	100 KCl 0.2 CaCl ₂ 2.8 MgCl ₂	100 RbCl 0.2 CaCl ₂ 2.8 MgCl ₂	100 CsCl 0.2 CaCl ₂ 2.8 MgCl ₂
5 mM HEPES, pH 7.4				

5 Reversal potentials were determined from tail currents after a 2 s depolarising pulse to +60 mV and corrected for liquid junction potentials that were determined experimentally. The permeability ratios were calculated according to $P_K / P_X = \exp (-F \cdot V_{rev} / R \cdot T)$.

To determine the voltage dependence of apparent open probability,
10 oocytes were clamped for 4 s to values between -80 mV to +50 mV in 10 mV steps, followed by a constant -30 mV test pulse. Tail currents extrapolated to $t=0$ were obtained from a mono-exponential fit, normalised to the value at 0 mV and used for the analysis of apparent p_{open} . Data analysis used PClamp6 and Microcal Origin 5.0.

Similar to KCNQ1, KCNQ2 and KCNQ3, also KCNQ4 yielded currents that
15 activated upon depolarisation (Fig. 1A). Compared to those other KCNQ channels, however, current activation was slower and occurred with a time constant in the order of 600 ms at +40 mV (KCNQ2/KCNQ3 channels have a corresponding time constant of ≈ 300 ms). This time constant was very sensitive to changes in temperature. Deactivation of currents at physiological resting potentials (≈ -70 mV) was considerably
20 faster (Fig. 1B). Similar to KCNQ2, macroscopic currents often showed some inward rectification at positive potentials. When oocytes were depolarised to +60 mV for 10 sec or more, an apparent slow inactivation of currents was observed that resembled the one described for KCNQ3. Currents began to activate at about -40 mV, with half-maximal activation at -10 mV (Fig. 2C). Ion substitution experiments showed that the
25 channel is highly selective for potassium (Fig. 1D). It has a $K^+ \approx Rb^+ > Cs^+ > Na^+$ permeability sequence. KCNQ4 currents were inhibited by more than 80% by 5 mM Ba⁺⁺.

We next examined the effect of the G285S mutation found in the affected family (Figs. 1E and 1F). The mutant channel did not yield any detectable currents in
30 the *Xenopus* oocyte expression system. KCNQ4_{G285S} was then injected at a 1:1 ratio with WT KCNQ4 to mimic the situation in a heterozygous DFNA2 patient. This

reduced currents by about 90%, indicating a strong dominant negative effect of the mutant. The degree of current reduction is compatible with the notion that the incorporation of one mutant subunit suffices to abolish the function of the tetrameric channel complex. The channels present in co-injected oocytes still showed a strong
 5 preference of potassium over sodium or calcium. This implies that the deafness is due to a quantitative loss of KCNQ4 potassium currents rather than to an influx of sodium or calcium.

KCNQ1 assembles with minK (IsK) to form channels that yield larger currents and activate much slower. We therefore tested by co-expression whether
 10 minK affects KCNQ4 as well. At concentrations (1 ng minK cRNA per oocyte) leading to drastic changes in KCNQ1 currents in parallel experiments, there was no significant change in KCNQ4 currents.

Different KCNQ subunits can form heteromeric channels. Co-expression of KCNQ2 with KCNQ3, but not with KCNQ1, gave currents that were about tenfold
 15 larger than those from homomeric channels. Since also KCNQ1 and KCNQ3 (and to a lesser degree also KCNQ2) are expressed in the cochlea, we investigated whether these proteins interact functionally. Oocytes co-injected (at the same total cRNA concentration) with KCNQ1 and KCNQ4 cRNAs yielded currents that seemed not different from a linear superposition of currents from the respective homomeric
 20 channels (Fig. 2A), and the same was true for oocytes co-expressing KCNQ2 and KCNQ4 (Fig. 2B). In addition, a dominant negative KCNQ1 mutant did not suppress KCNQ4 currents (Fig. 2A), and the same was true for the equivalent KCNQ2 mutant (Fig. 2B).

By contrast, co-expression of KCNQ3 with KCNQ4 yielded currents that
 25 were significantly larger than could be explained by a superposition of currents from the respective homomeric channels (Figs. 2C and 2D). Further, KCNQ4 currents were markedly suppressed by co-expressing a dominant negative KCNQ3 mutant (Fig. 2C). Importantly, currents from co-injected oocytes activated faster than KCNQ4 currents (Fig. 2D), and there was a ≈ 10 mV shift of the open probability towards negative
 30 voltages (Fig. 2E). Compared to KCNQ2/KCNQ3 channels, which may underlie the M-current, KCNQ3/KCNQ4 heteromers open at slightly more positive voltages. To compare KCNQ3/KCNQ4 channels to M-channels, we used the typical voltage-protocol employed for these channels and found currents superficially resembling M-currents (Fig. 2F). Linopirdine, a potent and rather specific inhibitor for M-currents,
 35 nearly completely inhibits KCNQ2/KCNQ3 channels at a concentration of 200 μ M. This concentration of Linopirdine inhibited KCNQ4 by about 30%, while a significantly larger inhibition ($\approx 75\%$) was observed with KCNQ3/KCNQ4 co-expression (Fig. 2G).

CLAIMS

1. An isolated polynucleotide having a nucleic acid sequence which is capable of hybridising under high stringency conditions with the polynucleotide sequence presented as SEQ ID NO: 1, its complementary strand, or a sub-sequence thereof.
2. The isolated polynucleotide according to claim 1, being at least 50% homologous, preferably more than 70%, more preferred more than 80%, even more preferred more than 90%, most preferred more than 95%, homologous to the polynucleotide sequence presented as SEQ ID NO: 1.
3. The isolated polynucleotide according to either of claims 1-2 being a cloned polynucleotide.
4. The isolated polynucleotide according to claim 3, in which the polynucleotide is cloned from, or produced on the basis of a cDNA library.
5. The isolated polynucleotide according to any of claims 1-4, comprising the polynucleotide sequence presented as SEQ ID NO: 1, or a sub-sequence hereof.
6. The isolated polynucleotide according to any of claims 1-4, comprising the polynucleotide sequence presented as SEQ ID NO: 1, or a sub-sequence hereof, including the mutation G935A.
7. The isolated polynucleotide according to any of claims 1-6, encoding a potassium channel, or a potassium channel subunit.
8. The isolated polynucleotide according to claim 7, encoding the KCNQ4 potassium channel subunit comprising the amino acid sequence represented by SEQ ID NO: 2, or a sub-sequence hereof.
9. The isolated polynucleotide according to claim 7, encoding a KCNQ4 variant, which variant has an amino acid sequence that has been changed by deletion of an amino acid residue, by insertion of an additional amino acid residue, or by substitution of an amino acid residue at one or more positions.

10. The isolated polynucleotide according to claim 9, which variant has an amino acid sequence that has been changed at one or more positions located in the conserved regions, as defined by Table 1.
- 5 11. The isolated polynucleotide according to claim 9, encoding the variant KCNQ4/G285S (i.e. KCNQ4/G333S according to the KCNQ1 numbering).
12. An isolated polynucleotide comprising any one of the sequences set forth in SEQ ID NOS: 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23,
10 24, 25, 26, 27, 28, 29, 30, 31, and 32.
13. A recombinantly produced polypeptide encoded by the polynucleotide according to claims 1-11.
- 15 14. The polypeptide according to claim 13, being a KCNQ4 potassium channel subunit comprising the amino acid sequence presented as SEQ ID No. 2.
15. The polypeptide according to claim 13, being a KCNQ4 variant, which variant has an amino acid sequence that has been changed by deletion of an amino acid residue, by insertion of an additional amino acid residue, or by substitution of an
20 amino acid residue at one or more positions.
16. The polypeptide according to claim 15, which variant has an amino acid sequence that has been changed at one or more positions located in the conserved regions, as defined by Table 1.
25
17. The polypeptide according to claim 15, being the variant KCNQ4/G285S (i.e. KCNQ4/G333S according to the KCNQ1 numbering).
- 30 18. A cell genetically manipulated by the incorporation of a heterologous polynucleotide according to any of claims 1-11.
19. The cell according to claim 18, genetically manipulated by the incorporation of a KCNQ4 channel subunit comprising the amino acid sequence presented as SEQ
35 ID NO: 2, or a sub-sequence hereof.
20. The cell according to claim 18, genetically manipulated by the incorporation of a KCNQ4 variant, which variant has an amino acid sequence that has been

changed by deletion of an amino acid residue, by insertion of an additional amino acid residue, or by substitution of an amino acid residue at one or more positions.

21. The cell according to claim 20, which variant has an amino acid sequence that has been changed at one or more positions located in the conserved regions, as defined by Table 1.
22. The cell according to claim 20, genetically manipulated by the incorporation of the variant KCNQ4/G285S (i.e. KCNQ4/G333S according to the KCNQ1 numbering).
23. The cell according to any of claims 18-22, genetically manipulated to co-express one or more KCNQ channel subunits.
24. The cell according to claim 23, genetically manipulated to co-express KCNQ4 and KCNQ1 channel subunits; KCNQ4 and KCNQ2 channel subunits; KCNQ4 and KCNQ3 channel subunits; KCNQ4 and KCNQ1 and KCNQ2 channel subunits; KCNQ4 and KCNQ1 and KCNQ3 channel subunits; KCNQ4 and KCNQ2 and KCNQ3 channel subunits; or KCNQ4 and KCNQ1 and KCNQ2 and KCNQ3 channel subunits.
25. The cell according to claim 23, genetically manipulated to co-express KCNQ3 and KCNQ4 channel subunits.
26. The cell according to any of claims 18-25, being an eukaryotic cell, in particular a mammalian cell, an oocyte, or a yeast cell.
27. The cell according to any claim 26, being a human embryonic kidney (HEK) cell, a HEK 293 cell, a BHK21 cell, a Chinese hamster ovary (CHO) cell, a *Xenopus laevis* oocyte (XLO) cell, a COS cell, or any other cell line able to express KCNQ potassium channels.
28. A membrane preparation derived from a cell according to any of claims 18-27.
29. A method for obtaining a substantially homogeneous source of a human potassium channel, comprising a KCNQ4 subunit, which method comprises the steps of culturing a cellular host having incorporated expressibly therein a

polynucleotide according to any of claims 1-11, and then recovering the cultured cells.

30. The method of claim 29, comprising the subsequent step of obtaining a membrane preparation from the cultured cells.
31. A method of screening a chemical compound for capability of binding to a potassium channel comprising at least one KCNQ4 channel subunit, which method comprises the steps of
 - (i) subjecting a KCNQ4 channel subunit containing cell according to claims 18-27, or a membrane preparation according to claim 28, to the action of a KCNQ4 binding agent to form a complex with the KCNQ4 channel subunit containing cell;
 - (ii) subjecting the complex of step (i) to the action of the chemical compound to be tested; and
 - (iii) detecting the displacement of the KCNQ4 binding agent from the complex with the KCNQ4 channel subunit containing cell or membrane preparation.
32. The method of claim 31, wherein the KCNQ4 channel subunit containing cell is a cell according to any of claims 18-27, or a membrane preparation according to claim 28.
33. The method of either of claims 31-32, in which the KCNQ4 binding agent is
 - radioactively labelled 1,3-dihydro-1-phenyl-3,3-bis(4-pyridylmethyl)-2H-indol-2-one (Linopirdine); or
 - radioactively labelled 10,10-bis(4-pyridinyl-methyl)-9(10H)-anthracenone.
34. The method of claim 33, which compounds have been marked with ^3H .
35. The method of either of claims 33-34, wherein the displacement of the KCNQ4 binding agent from the complex with the KCNQ4 channel subunit containing cell is detected by measuring the amount of radioactivity by conventional liquid scintillation counting.
36. A method of screening a chemical compound for activity on a potassium channel comprising at least one KCNQ4 channel subunit, which method comprises the steps of

- (i) subjecting a KCNQ4 channel subunit containing cell to the action of the chemical compound; and
- (ii) monitoring the membrane potential, the current, the potassium flux, or the secondary calcium influx of the KCNQ4 channel subunit containing cell.

5

37. The method of claim 36, wherein the KCNQ4 channel subunit containing cell is a cell according to any of claims 18-27.

10

38. The method of either of claims 36-37, wherein monitoring of the membrane potential of the KCNQ4 channel subunit containing cell is performed by patch clamp techniques.

15

39. The method of either of claims 36-37, wherein monitoring of the membrane potential of the KCNQ4 channel subunit containing cell is performed using fluorescence methods.

20

40. A chemical compound identified by the method of claims 31-35, and/or by the method of claims 36-39.

25

41. Use of the chemical compound according to claim 40 for diagnosis, treatment, prevention or alleviation of diseases related to tinnitus, loss of hearing, in particular progressive hearing loss, neonatal deafness, and presbycusis (deafness of the elderly); and diseases or adverse conditions of the CNS, including affective disorders, Alzheimer's disease, anxiety, ataxia, CNS damage caused by trauma, stroke or neurodegenerative illness, cognitive deficits, compulsive behaviour, dementia, depression, Huntington's disease, mania, memory impairment, memory disorders, memory dysfunction, motion disorders, motor disorders, neurodegenerative diseases, Parkinson's disease and Parkinson-like motor disorders, phobias, Pick's disease, psychosis, schizophrenia, spinal cord damage, stroke, and tremor.

30

42. The use according to claim 41, wherein the chemical compound is
 1,3-dihydro-1-phenyl-3,3-bis(4-pyridylmethyl)-2H-indol-2-one (Linopirdine);
 or
 10,10-bis(4-pyridinyl-methyl)-9(10H)-anthracenone.

35

43. Use of a polynucleotide sequence according to any of claims 1-12, for the screening of genetic materials for individuals having this mutations.

44. A transgenic animal comprising a knock-out mutation of the endogenous *KCNQ4* gene, a mutated *KCNQ4* gene, or genetically manipulated in order to over-express the *KCNQ4* gene or to over-express mutated *KCNQ4* gene.

5

45. The transgenic animal according to claim 44, being a knock-out animal in which the gene is totally deleted in a homozygous state.

46. The transgenic animal according to claim 44, comprising a mutated *KCNQ4* gene.

10

47. The transgenic animal according to any of claims 44-46, being a transgenic rodent, in particular a hamster, a guinea pig, a rabbit, or a rat, a transgenic pig, a transgenic cattle, a transgenic sheep, or a transgenic goat.

15

48. Use of the transgenic animal according to any of claims 44-47 for the *in vivo* screening of therapeutic compounds.

49. The use according to claim 48, for the screening of drugs affecting diseases or conditions associated with hearing loss or tinnitus.

20

50. A method for the identification, localization, isolation or amplification a polynucleotide according to any one of claims 1-11, comprising using a polynucleotide according to claim 12 as a primer or a probe.

25

51. An antibody capable of binding one or more polypeptides as claimed in any one of claims 13-17.

52. The antibody of claim 51 being a monoclonal antibody.

**TITLE: NOVEL POTASSIUM CHANNELS AND
GENES ENCODING THESE POTASSIUM CHANNELS**

ABSTRACT

This invention relates to novel potassium channels and genes encoding these channels. More specifically the invention provides isolated polynucleotides encoding the KCNQ4 potassium channel, cells transformed with these polynucleotides, transgenic animals comprising genetic mutations, and the use of the transformed cells and the transgenic animals for the *in vitro* and *in vivo* screening of drugs affecting KCNQ4 containing potassium channels.

1/4




Fig. 1A-1C

2/4




Fig. 1D-1F

3/4




Fig. 2A-2C




Fig. 2D-2G

<120> Novel potassium channels and genes encoding these potassium channels.

<140>

<160> 32

 $\langle 210 \rangle$ 1

<211> 2335

<212> DNA

<213> Homo sapiens

 $\langle 220 \rangle$

<221> gene

 $\langle 222 \rangle \quad (1) \dots (2335)$ $\langle 220 \rangle$

<221> CDS

<222> (83) .. (2170)

<223> KCNQ4

<400> 1

agccatgcgt ctctgagcgc cccgagcgcg ccccgcccc ggaccgtgcc cgggccccgg 60

cgccccagc ccggcgcgc cc atg gcc gag gcc ccc ccg cgc cgc ctc ggc 112
Met Ala Glu Ala Pro Pro Arg Arg Leu Gly
1 5 10

ctg ggt ccc ccg ccc ggg gac gcc ccc cgc gcg gag cta gtg gcg ctc 160
Leu Gly Pro Pro Pro Gly Asp Ala Pro Arg Ala Glu Leu Val Ala Leu
15 20 25

acg gcc gtg cag agc gaa cag ggc gag gcg ggc ggg ggc ggc tcc ccg 208
Thr Ala Val Gln Ser Glu Gln Gly Glu Ala Gly Gly Gly Gly Ser Pro
30 35 40

cgc cgc ctc ggc ctc ctg ggc agc ccc ctg ccg ccg ggc gcg ccc ctc 256
Arg Arg Leu Gly Leu Leu Gly Ser Pro Leu Pro Pro Gly Ala Pro Leu
45 50 55

cct ggg ccg ggc tcc ggc tcg ggc tcc gcc tgc ggc cag cgc tcc tcg 304
Pro Gly Pro Gly Ser Gly Ser Gly Ser Ala Cys Gly Gln Arg Ser Ser
60 65 70

gcc gcg cac aag cgc tac cgc cgc ctg cag aac tgg gtc tac aac gtg 352
Ala Ala His Lys Arg Tyr Arg Arg Leu Gln Asn Trp Val Tyr Asn Val
75 80 85 90

ctg	gag	cgg	ccc	cgc	ggc	tgg	gcc	ttc	gtc	tac	cac	gtc	ttc	ata	ttt	400
Leu	Glu	Arg	Pro	Arg	Gly	Trp	Ala	Phe	Val	Tyr	His	Val	Phe	Ile	Phe	
				95					100					105		

tgt	ctg	gtc	ttc	agc	tgc	ctg	gtg	ctg	tct	gtg	ctg	tcc	act	atc	cag	448
Leu	Leu	Val	Phe	Ser	Cys	Leu	Val	Leu	Ser	Val	Leu	Ser	Thr	Ile	Gln	
			110					115					120			

2

gag	cac	cag	gaa	ctt	gcc	aac	gag	tgt	ctc	ctc	atc	ttg	gaa	ttc	gtg	496
Glu	His	Gln	Glu	Leu	Ala	Asn	Glu	Cys	Leu	Leu	Ile	Leu	Glu	Phe	Val	
		125					130					135				
atg	atc	gtg	gtt	ttc	ggc	ttg	gag	tac	atc	gtc	cgg	gtc	tgg	tcc	gcc	544
Met	Ile	Val	Val	Phe	Gly	Leu	Glu	Tyr	Ile	Val	Arg	Val	Trp	Ser	Ala	
	140					145					150					
gga	tgc	tgc	tgc	cgc	tac	cga	gga	tgg	cag	ggt	cgc	ttc	cgc	ttt	gcc	592
Gly	Cys	Cys	Cys	Arg	Tyr	Arg	Gly	Trp	Gln	Gly	Arg	Phe	Arg	Phe	Ala	
155					160					165					170	
aga	aag	ccc	ttc	tgt	gtc	atc	gac	ttc	atc	gtg	ttc	gtg	gcc	tgc	gtg	640
Arg	Lys	Pro	Phe	Cys	Val	Ile	Asp	Phe	Ile	Val	Phe	Val	Ala	Ser	Val	
			175						180					185		
gcc	gtc	atc	gcc	gcg	ggt	acc	cag	ggc	aac	atc	ttc	gcc	acg	tcc	gcg	688
Ala	Val	Ile	Ala	Ala	Gly	Thr	Gln	Gly	Asn	Ile	Phe	Ala	Thr	Ser	Ala	
			190					195					200			
ctg	cgc	agc	atg	cgc	ttc	ctg	cag	atc	ctg	cgc	atg	gtg	cgc	atg	gac	736
Leu	Arg	Ser	Met	Arg	Phe	Leu	Gln	Ile	Leu	Arg	Met	Val	Arg	Met	Asp	
		205					210					215				
cgc	cgc	ggc	ggc	acc	tgg	aag	ctg	ctg	ggc	tca	gtg	gtc	tac	gcg	cat	784
Arg	Arg	Gly	Gly	Thr	Trp	Lys	Leu	Leu	Gly	Ser	Val	Val	Tyr	Ala	His	
		220				225					230					
agc	aag	gag	ctg	atc	acc	gcc	tgg	tac	atc	ggg	ttc	ctg	gtg	ctc	atc	832
Ser	Lys	Glu	Leu	Ile	Thr	Ala	Trp	Tyr	Ile	Gly	Phe	Leu	Val	Leu	Ile	
235					240					245					250	
ttc	gcc	tcc	ttc	ctg	gtc	tac	ctg	gcc	gag	aag	gac	gcc	aac	tcc	gac	880
Phe	Ala	Ser	Phe	Leu	Val	Tyr	Leu	Ala	Glu	Lys	Asp	Ala	Asn	Ser	Asp	
			255						260					265		
ttc	tcc	tcc	tac	gcc	gac	tgc	ctc	tgg	tgg	ggg	acg	att	aca	ttg	aca	928
Phe	Ser	Ser	Tyr	Ala	Asp	Ser	Leu	Trp	Trp	Gly	Thr	Ile	Thr	Leu	Thr	
			270					275					280			
acc	atc	ggc	tat	ggt	gac	aag	aca	cgc	cac	aca	tgg	ctg	ggc	agg	gtc	976
Thr	Ile	Gly	Tyr	Gly	Asp	Lys	Thr	Pro	His	Thr	Trp	Leu	Gly	Arg	Val	
		285					290					295				
ctg	gct	gct	ggc	ttc	gcc	tta	ctg	ggc	atc	tct	ttc	ttt	gcc	ctg	cct	1024
Leu	Ala	Ala	Gly	Phe	Ala	Leu	Leu	Gly	Ile	Ser	Phe	Phe	Ala	Leu	Pro	
	300					305					310					
gcc	ggc	atc	cta	ggc	tcc	ggc	ttt	gcc	ctg	aag	gtc	cag	gag	cag	cac	1072
Ala	Gly	Ile	Leu	Gly	Ser	Gly	Phe	Ala	Leu	Lys	Val	Gln	Glu	Gln	His	
315					320					325					330	
cgg	cag	aag	cac	ttc	gag	aag	cgg	agg	atg	cgc	gca	gcc	aac	ctc	atc	1120
Arg	Gln	Lys	His	Phe	Glu	Lys	Arg	Arg	Met	Pro	Ala	Ala	Asn	Leu	Ile	
			335						340					345		
cag	gct	gcc	tgg	cgc	ctg	tac	tcc	acc	gat	atg	agc	cgg	gcc	tac	ctg	1168
Gln	Ala	Ala	Trp	Arg	Leu	Tyr	Ser	Thr	Asp	Met	Ser	Arg	Ala	Tyr	Leu	
		350						355					360			
aca	gcc	acc	tgg	tac	tac	tat	gac	agt	atc	ctc	cca	tcc	ttc	aga	gag	1216
Thr	Ala	Thr	Trp	Tyr	Tyr	Tyr	Asp	Ser	Ile	Leu	Pro	Ser	Phe	Arg	Glu	
		365					370					375				

ctg gcc ctc ttg ttt gag cac gtg caa cgg gcc cgc aat ggg ggc cta	1264
Leu Ala Leu Leu Phe Glu His Val Gln Arg Ala Arg Asn Gly Gly Leu	
380 385 390	
cgg ccc ctg gag gtg cgg cgg gcg cgg gta ccc gac gga gca ccc tcc	1312
Arg Pro Leu Glu Val Arg Arg Ala Pro Val Pro Asp Gly Ala Pro Ser	
395 400 405 410	
cgt tac ccg ccc gtt gcc acc tgc cac cgg ccg ggc agc acc tcc ttc	1360
Arg Tyr Pro Pro Val Ala Thr Cys His Arg Pro Gly Ser Thr Ser Phe	
415 420 425	
tgc cct ggg gaa agc agc cgg atg ggc atc aaa gac cgc atc cgc atg	1408
Cys Pro Gly Glu Ser Ser Arg Met Gly Ile Lys Asp Arg Ile Arg Met	
430 435 440	
ggc agc tcc cag cgg cgg acg ggt cct tcc aag cag cag ctg gca cct	1456
Gly Ser Ser Gln Arg Arg Thr Gly Pro Ser Lys Gln Gln Leu Ala Pro	
445 450 455	
cca aca atg ccc acc tcc cca agc agc gag cag gtg ggt gag gcc acc	1504
Pro Thr Met Pro Thr Ser Pro Ser Ser Glu Gln Val Gly Glu Ala Thr	
460 465 470	
agc ccc acc aag gtg caa aag agc tgg agc ttc aat gac cgc acc cgc	1552
Ser Pro Thr Lys Val Gln Lys Ser Trp Ser Phe Asn Asp Arg Thr Arg	
475 480 485 490	
ttc cgg gca tct ctg aga ctc aaa ccc cgc acc tct gct gag gat gcc	1600
Phe Arg Ala Ser Leu Arg Leu Lys Pro Arg Thr Ser Ala Glu Asp Ala	
495 500 505	
ccc tca gag gaa gta gca gag gag aag agc tac cag tgt gag ctc acg	1648
Pro Ser Glu Glu Val Ala Glu Glu Lys Ser Tyr Gln Cys Glu Leu Thr	
510 515 520	
gtg gac gac atc atg cct gct gtg aag aca gtc atc cgc tcc atc agg	1696
Val Asp Asp Ile Met Pro Ala Val Lys Thr Val Ile Arg Ser Ile Arg	
525 530 535	
att ctc aag ttc ctg gtg gcc aaa agg aaa ttc aag gag aca ctg cga	1744
Ile Leu Lys Phe Leu Val Ala Lys Arg Lys Phe Lys Glu Thr Leu Arg	
540 545 550	
ccg tac gac gtg aag gac gtc att gag cag tac tca gca ggc cac ctg	1792
Pro Tyr Asp Val Lys Asp Val Ile Glu Gln Tyr Ser Ala Gly His Leu	
555 560 565 570	
gac atg ctg ggc cgg atc aag agc ctg caa act cgg gtg gac caa att	1840
Asp Met Leu Gly Arg Ile Lys Ser Leu Gln Thr Arg Val Asp Gln Ile	
575 580 585	
gtg ggt cgg ggg ccc ggg gac agg aag gcc cgg gag aag ggc gac aag	1888
Val Gly Arg Gly Pro Gly Asp Arg Lys Ala Arg Glu Lys Gly Asp Lys	
590 595 600	
ggg ccc tcc gac gcg gag gtg gtg gat gaa atc agc atg atg gga cgc	1936
Gly Pro Ser Asp Ala Glu Val Val Asp Glu Ile Ser Met Met Gly Arg	
605 610 615	
gtg gtc aag gtg gag aag cag gtg cag tcc atc gag cac aag ctg gac	1984
Val Val Lys Val Glu Lys Gln Val Gln Ser Ile Glu His Lys Leu Asp	
620 625 630	

4

ctg ctg ttg ggc ttc tat tgc cgc tgc ctg cgc tct ggc acc tgc gcc 2032
 Leu Leu Leu Gly Phe Tyr Ser Arg Cys Leu Arg Ser Gly Thr Ser Ala
 635 640 645 650
 agc ctg ggc gcc gtg caa gtg ccg ctg ttc gac ccc gac atc acc tcc 2080
 Ser Leu Gly Ala Val Gln Val Pro Leu Phe Asp Pro Asp Ile Thr Ser
 655 660 665
 gac tac cac agc cct gtg gac cac gag gac atc tcc gtc tcc gca cag 2128
 Asp Tyr His Ser Pro Val Asp His Glu Asp Ile Ser Val Ser Ala Gln
 670 675 680
 acg ctc agc atc tcc cgc tgc gtc agc acc aac atg gac tga 2170
 Thr Leu Ser Ile Ser Arg Ser Val Ser Thr Asn Met Asp
 685 690 695
 gggacttctc agaggcaggg cagcacacgg ccagccccgc ggcttgggcg tccgactgcc 2230
 ctctgaggcc tccggactcc tctcgtactt gaactcactc cctcacgggg agagagacca 2290
 caccgagtat tgagctgcct gaggggcggt ggtacctgct gtggg 2335

<210> 2
 <211> 695
 <212> PRT
 <213> Homo sapiens

<400> 2
 Met Ala Glu Ala Pro Pro Arg Arg Leu Gly Leu Gly Pro Pro Pro Gly
 1 5 10 15
 Asp Ala Pro Arg Ala Glu Leu Val Ala Leu Thr Ala Val Gln Ser Glu
 20 25 30
 Gln Gly Glu Ala Gly Gly Gly Gly Ser Pro Arg Arg Leu Gly Leu Leu
 35 40 45
 Gly Ser Pro Leu Pro Pro Gly Ala Pro Leu Pro Gly Pro Gly Ser Gly
 50 55 60
 Ser Gly Ser Ala Cys Gly Gln Arg Ser Ser Ala Ala His Lys Arg Tyr
 65 70 75 80
 Arg Arg Leu Gln Asn Trp Val Tyr Asn Val Leu Glu Arg Pro Arg Gly
 85 90 95
 Trp Ala Phe Val Tyr His Val Phe Ile Phe Leu Leu Val Phe Ser Cys
 100 105 110
 Leu Val Leu Ser Val Leu Ser Thr Ile Gln Glu His Gln Glu Leu Ala
 115 120 125
 Asn Glu Cys Leu Leu Ile Leu Glu Phe Val Met Ile Val Val Phe Gly
 130 135 140
 Leu Glu Tyr Ile Val Arg Val Trp Ser Ala Gly Cys Cys Cys Arg Tyr
 145 150 155 160
 Arg Gly Trp Gln Gly Arg Phe Arg Phe Ala Arg Lys Pro Phe Cys Val
 165 170 175
 Ile Asp Phe Ile Val Phe Val Ala Ser Val Ala Val Ile Ala Ala Gly
 180 185 190

5

Thr Gln Gly Asn Ile Phe Ala Thr Ser Ala Leu Arg Ser Met Arg Phe
 195 200 205
 Leu Gln Ile Leu Arg Met Val Arg Met Asp Arg Arg Gly Gly Thr Trp
 210 215 220
 Lys Leu Leu Gly Ser Val Val Tyr Ala His Ser Lys Glu Leu Ile Thr
 225 230 235 240
 Ala Trp Tyr Ile Gly Phe Leu Val Leu Ile Phe Ala Ser Phe Leu Val
 245 250 255
 Tyr Leu Ala Glu Lys Asp Ala Asn Ser Asp Phe Ser Ser Tyr Ala Asp
 260 265 270
 Ser Leu Trp Trp Gly Thr Ile Thr Leu Thr Thr Ile Gly Tyr Gly Asp
 275 280 285
 Lys Thr Pro His Thr Trp Leu Gly Arg Val Leu Ala Ala Gly Phe Ala
 290 295 300
 Leu Leu Gly Ile Ser Phe Phe Ala Leu Pro Ala Gly Ile Leu Gly Ser
 305 310 315 320
 Gly Phe Ala Leu Lys Val Gln Glu Gln His Arg Gln Lys His Phe Glu
 325 330 335
 Lys Arg Arg Met Pro Ala Ala Asn Leu Ile Gln Ala Ala Trp Arg Leu
 340 345 350
 Tyr Ser Thr Asp Met Ser Arg Ala Tyr Leu Thr Ala Thr Trp Tyr Tyr
 355 360 365
 Tyr Asp Ser Ile Leu Pro Ser Phe Arg Glu Leu Ala Leu Leu Phe Glu
 370 375 380
 His Val Gln Arg Ala Arg Asn Gly Gly Leu Arg Pro Leu Glu Val Arg
 385 390 395 400
 Arg Ala Pro Val Pro Asp Gly Ala Pro Ser Arg Tyr Pro Pro Val Ala
 405 410 415
 Thr Cys His Arg Pro Gly Ser Thr Ser Phe Cys Pro Gly Glu Ser Ser
 420 425 430
 Arg Met Gly Ile Lys Asp Arg Ile Arg Met Gly Ser Ser Gln Arg Arg
 435 440 445
 Thr Gly Pro Ser Lys Gln Gln Leu Ala Pro Pro Thr Met Pro Thr Ser
 450 455 460
 Pro Ser Ser Glu Gln Val Gly Glu Ala Thr Ser Pro Thr Lys Val Gln
 465 470 475 480
 Lys Ser Trp Ser Phe Asn Asp Arg Thr Arg Phe Arg Ala Ser Leu Arg
 485 490 495
 Leu Lys Pro Arg Thr Ser Ala Glu Asp Ala Pro Ser Glu Glu Val Ala
 500 505 510
 Glu Glu Lys Ser Tyr Gln Cys Glu Leu Thr Val Asp Asp Ile Met Pro
 515 520 525
 Ala Val Lys Thr Val Ile Arg Ser Ile Arg Ile Leu Lys Phe Leu Val
 530 535 540

6

Ala Lys Arg Lys Phe Lys Glu Thr Leu Arg Pro Tyr Asp Val Lys Asp
 545 550 555 560

Val Ile Glu Gln Tyr Ser Ala Gly His Leu Asp Met Leu Gly Arg Ile
 565 570 575

Lys Ser Leu Gln Thr Arg Val Asp Gln Ile Val Gly Arg Gly Pro Gly
 580 585 590

Asp Arg Lys Ala Arg Glu Lys Gly Asp Lys Gly Pro Ser Asp Ala Glu
 595 600 605

Val Val Asp Glu Ile Ser Met Met Gly Arg Val Val Lys Val Glu Lys
 610 615 620

Gln Val Gln Ser Ile Glu His Lys Leu Asp Leu Leu Leu Gly Phe Tyr
 625 630 635 640

Ser Arg Cys Leu Arg Ser Gly Thr Ser Ala Ser Leu Gly Ala Val Gln
 645 650 655

Val Pro Leu Phe Asp Pro Asp Ile Thr Ser Asp Tyr His Ser Pro Val
 660 665 670

Asp His Glu Asp Ile Ser Val Ser Ala Gln Thr Leu Ser Ile Ser Arg
 675 680 685

Ser Val Ser Thr Asn Met Asp
 690 695

<210> 3
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 3
 catgctgtctc tgagcgcccc gagg

24

<210> 4
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 4
 aggccaggct tgcgcgggga aacg

24

<210> 5
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 5
 cagcacagag ctgtaactcc agg

23

<210> 6
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 6
 aagctgctct ctgagccatg g 21

<210> 7
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 7
 gctgggtccg cgctgtgacc 20

<210> 8
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 8
 ggtctccagg gtcagagtcg 20

<210> 9
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 9
 tccgggtccg tgcgcggggt a 21

<210> 10
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 10
 gagacagccc ctctgacctc g 21

<210> 11
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 11
atccctttcc cgtgtggaag c 21

<210> 12
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 12
agtcacgatg ggcagacctc g 21

<210> 13
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 13
cctcatgatc aggtcctac c 21

<210> 14
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 14
atgtgtgaca ggggtgagc 19

<210> 15
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 15
aaggatgggg acacccttgc 20

<210> 16
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 16
acacaggggtt gacacacc 18

<210> 17
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 17
 gctctgggta acccacaact g 21

<210> 18
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 18
 gctccccctgg gagccatcac c 21

<210> 19
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 19
 tgagctcagg agctctgtgc 20

<210> 20
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 20
 acccacgaag tggctgaagg c 21

<210> 21
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 21
 gtcctaagtc agctttgtcc 20

<210> 22
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

 <400> 22
 cctcagccgg ccctcgatcg 20

 <210> 23
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: PCR primer

 <400> 23
 cactctactg gtggtttggc 20

 <210> 24
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: PCR primer

 <400> 24
 ctcctgacct caagtgatcc 20

 <210> 25
 <211> 21
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: PCR primer

 <400> 25
 gatagcaaag agatggagag g 21

 <210> 26
 <211> 21
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: PCR primer

 <400> 26
 aactcagctg cagcagtgag c 21

 <210> 27
 <211> 21
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: PCR primer

 <400> 27
 gtgccttctc cttcatcagg c 21

<210> 28
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 28
aacgcatacct ccccatgtca

20

<210> 29
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 29
tttgtgcttc ccagataagc

20

<210> 30
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 30
cgtgagggag tgagttcaag tacg

24

<210> 31
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 31
agtacctgat ggagcgcct ctcg

24

<210> 32
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 32
tcattccaccg taagtcaca ctgg

24